

Fossilium Catalogus

II: Plantae.

Editus a

W. Jongmans.

Pars 17:

W. N. Edwards

Dicotyledones (Ligna).



W. Junk
Berlin W. 15
26. I. 1931

Ds

115

Do 115, v,

W. JUNK, Verlag und Antiquariat für Naturwissenschaften
Berlin, W. 15

Fossilium Catalogus

I: Animalia :

- Pars 1: F. Frech, Ammoneae devonicae [Clymeniidae, Aphyllitidae, Gephyroceratidae, Cheiloceratidae]. 1913. 42 p.
2, 15: W. v. Teppner, Lamellibranchiata tertiaria. 'Anisomyaria' I. II. 1914—1922. 296 p.
3: Ch. Schuchert, Stellerioidea palaeozoica. 1914. 53 p.
4: F. de Huene, Saurischia et Ornithischia triadica ['Dinosauria triadica']. 1914. 21 p.
5: J. Felix, Anthozoa palaeocretacea. 1914. 84 p.
6: J. Felix, Anthozoa cenomanica. 1914. 59 p.
7: J. Felix, Anthozoa neocretacea. 1914. 129 p.
8: C. Diener, Cephalopoda triadica. 1915. 369 p.
9: E. Hennig, Stegosauria. 1915. 16 p.
10: C. Diener, Brachiopoda triadica. 1920. 109 p.
11: C. v. Buelow, Cephalopoda dibranchiata. 1920. 313 p.
12: K. Lambrecht, Aves. 1921. 108 p.
13: C. Diener, Cnidaria triadica. 1921. 46 p.
14: C. Diener, Ammonoidea permiana. 1921. 36 p.
16: A. Handlirsch, Insecta palaeozoica. 1922. 230 p.
17, 18, 20—23, 32, 38, 40, 43, 46: W. Wenz, Gastropoda extramarina tertiaria. 11 partes. 1923—1930. 3387 p.
19: C. Diener, Lamellibranchiata triadica. 1923. 260 p.
24: O. Posthumus, Otolithi Piscium. 1924. 42 p.
25: C. Diener, Eurypterida. 1924. 29 p.
26: A. Kieslinger, Medusae fossiles. 1924. 20 p.
27: F. v. Nopcsa, Osteologia Reptilium fossilium et recentium. 1926. 391 p.
28: J. Felix, Anthozoa eocaenica et oligocaenica. 1925. 296 p.
29: C. Diener, Ammonoidea neocretacea. 1925. 244 p.
30: W. Deecke, Trigoniidae mesozoicae (Myophoria exclusa). 1925. 306 p.
31: W. O. Dietrich, Gastropoda mesozoica: Fam. Nerineidae. 1925. 166 p.
33: W. Deecke, Pisces triadici. 1926. 203 p.
34: C. Diener, Glossophora triadica. 1926. 244 p.
35: J. Felix, Anthozoa miocaenica. 1927. 194 p.
36: O. Kühn, Hydrozoa. 1928. 116 p.
37: Rud, et E. Richter, Trilobitae neodevonici. 1928. 162 p.
39: W. Deecke, Echinoidea jurassica. 1929. 540 p.
41: M. F. Glaessner, Crustacea decapoda. 1929. 464 p.

Fortsetzung siehe 3. u. 4. Umschlagseite.

Fossilium Catalogus

II: Plantae.

Editus a

W. Jongmans.

Pars 17:

W. N. Edwards


Dicotyledones (Ligna).



~~Wpisano do inwentarza
ZAKŁADU GEOLOGII~~

~~Dział B Nr 65
Dnia 9.10. 1946.~~

W. Junk
Berlin W. 15
1931.



3644

0

Contents.

	Page
Introduction	3
Bibliography	6
Alphabetical Catalogue of Genera and Species	16
Systematic List of Families and Genera	83
Geographical Index	85
Addenda	90
Index of Names	91



1931.254



Introduction.

No general review of the literature of structurally preserved dicotyledonous woods has appeared since 1890, when Kaiser listed, with very few omissions, the species described up to that date. The present catalogue, although a slight departure from the strictly systematic treatment of dicotyledonous families in the Fossilium Catalogus, may therefore be of use to workers in this field.

The genera and species of woods are arranged in alphabetical order, and a systematic index under families is given at the end. The bibliographic references are set out as in Fossilium Catalogus, Pars 14, Sapindaceae. When two or more papers by the same author have been published in one year, they are distinguished by the addition of letters thus; 1870, 1870a, 1870b.

Synonyms, nomina nuda, rejected names, and misidentifications are enclosed in square brackets, and full synonyms usually appear only under what is regarded as the correct, or the best established name. Cross-references are always given.

In general, records of woods in Pleistocene and Holocene deposits have only been included when there is discussion or illustration of the internal structure.

In some instances it has been possible to include in the remarks an indication of the whereabouts of the type material, but in the time at my disposal I have been unable to confirm all such references, nor to extend enquiries to cases where the original authors are silent on this matter.

NOMENCLATURE. Form-genera of fossil woods are usually made by adding the termination **-oxylon** to the name of the recent genus or family to which the wood is presumed to belong. Many of the older names, and especially Unger's, have the termination **-inium**, and several of these, such as **Quercinium**, are well-established and in general use. Others, such as **Betulinium**, have by certain authors been replaced by names like **Betuloxylon**. This seems to me quite unnecessary, and although there is a strong case for not submitting form-genera strictly to the rules of priority, I see no reason for giving up all the older generic names ending in **-inium**. As a matter of mere convenience, it may be pointed out that to substitute names ending in **-oxylon** for names ending in **-inium** would involve far more changes than the opposite course of retaining the older form-genera. Gothan argues that the **-oxylon** ending at once indicates the nature of the fossil, but so also, and perhaps more exclusively, does the **-inium** ending. It might happen that one wished to propose a form-genus for woods presumed to belong to the **Erythroxylaceae**, or the genera **Haematoxylon** or **Sideroxylon**, and the **-inium** termination would in such cases be much more appropriate.

Pleistocene and Holocene woods, the great majority of which certainly belong to living genera, and probably to living species, should be recorded under recent generic names. Nothing is gained by calling a Pleistocene elm wood **Ulmoxylon** rather than **Ulmus** sp., and to iden-

tify a comparatively recent willow-like wood with *Salicinoxyloxyon miocenicum*, as Fietz has done (1926), may cause confusion and is to be deprecated. The facts of the case are sufficiently, and indeed more clearly, indicated by recording such wood as *Salix* ? sp., as Fietz himself did in a similar case (1926a). Many of the geologically older woods may of course also belong to recent genera, but the element of uncertainty is greater.

Caspary's use of recent generic names for fossil woods which were often of uncertain generic position cannot be defended; his arguments concerning the nomenclature of fossil woods have been effectively criticized by Felix (1894), who was also able to show that a wood described by Caspary as *Schinus* was much more probably a member of the *Rhamnaceae*, and therefore was more appropriately to be termed *Rhamnacinium*.

For other nomenclatorial vagaries see the remarks below on *Alnoxyloxyon*, *Betulinium priscum*, *Betulinium* sp., *Fraxinus excelsior*, *Plataninium* sp., and *Salicinium antiquum*.

There are numerous dicotyledonous woods which cannot at present be placed even in families. The situation is, in fact, much as it was when Brongniart wrote in 1822. He proposed the name *Exogenites* for dicotyledonous woods in general, but this name is open to objection, and especially because dicotyledons, or exogens, included in those days such diverse groups as conifers, cycads, and some lycopods. I therefore suggest that Schleiden's name *Dryoxyloxyon*, expressly founded for the wood of broad-leaved trees (*Laubhölzer*) should be revived as a useful form-genus for dicotyledonous woods of uncertain position.

The practise of instituting entirely new genera (such as *Bronnites*, *Petzholdia*), started by Unger, has unfortunately continued to the present day (e. g. *Aulacoxylon*, *Sumatroxyloxyon*). I have transferred a few of Unger's genera to *Dryoxyloxyon*, when his name was pre-occupied by a living plant, but, failing a re-examination of the types, it does not seem worth while to rename the mass of species inadequately described and figured by Unger. All species of the following genera (very few of which have been figured at all) must be regarded as unidentifiable unless Unger's slides can be found and examined: *Brongniartites*, *Bronnites*, *Cottaites*, *Fichtelites*, *Fraasia*, *Klippsteinia*, *Meyenites*, *Mirbellites*, *Mohlites*, *Petzholdia*, *Piccolominites*, *Pritchardia*, *Rosthornia*, *Schleidenites*, *Sillimannia*, *Tchihatcheffites*, and *Withamia*. The following species of Unger's are also not figured and inadequately described: *Juglandinium mediterraneum*, *Laurinium guatemalense*, *L. xyloides*, *Rhodium juglandinum*, and *Salicinium populinum*.

Other genera of uncertain affinities (not always adequately described) which might well have been included in *Dryoxyloxyon* are: *Aachenoxylon*, *Cantia*, *Hythia*, *Jugloxylon*, *Palackya*, *Pataloxylon*, *Populocaulis*, *Sabulia*, *Sjögrenia*, *Stephanoxylon*, *Suevioxyloxyon*, *Sumatroxyloxyon*, *Trochodendromagnolia*, and *Zittelia*. It is very doubtful whether the multiplication of such names serves any useful purpose, and if it is really necessary to name unidentified dicotyledons I would strongly urge the use of the designation *Dryoxyloxyon*.

IDENTIFICATION. The possibility of identifying fossil dicotyledonous woods depends on the state of our knowledge of recent woody structure. A few references to works dealing with recent woods have been included in the bibliography. Solereder (1899, 1908) is the principal source of information on the anatomy of dicotyledonous families. For European woods see Muller (1888), Tetzlaff (1881), Chalk & Rendle (1929). For Eastern Asiatic woods see Foxworthy (1907, 1909), Kanehira (1921, 1921a), Moll & Janssonius (1906). Other useful works are Hopkinson (1912), Houlbert (1893), Moeller (1876), Stone (1904, 1921), and Wilhelm (1903). Nordlinger's "Holzquerschnitte" (1852) should also be mentioned, and also two sets of photographs of wood sections: James

Weale's Microphotographs of Woods, of which there are 1200 or more, and R. B. Hough's American Woods, of which 14 parts have appeared between 1888 and 1928, dealing with 350 woods.

For discussions of the problems connected with the identification of fossil dicotyledonous woods see among others Bailey (1924), Felix (1883a), Kräusel (1922), and Moll & Janssonius (1912).

METHODS OF PREPARATION and treatment of lignite, brown coal, charcoal, and so on are described by Bauer (1927), Fietz (1926, 1926a), Wittmack & Buchwald (1902).

PETRIFICATION AND MODES OF OCCURRENCE. The following papers may be mentioned: Barron (1905), Becke (1912), Chhibber (1928), Donath (1928), Johnson (1876), Kuntze (1895), Schweinfurth (1882), Stokes (1837) and Udluft (1926). The whole question of how plants become petrified is reviewed by St. John (1927), who gives numerous references to the earlier literature.

GEOGRAPHICAL DISTRIBUTION AND AGE. The systematic index under families is followed by a geographical index showing the distribution of described fossil woods. Numerous brief records of the occurrence of 'fossil wood' or 'dicotyledonous wood' without any further particulars, are scattered through the geological literature; they have been omitted from this catalogue.

The great majority of the species listed are of Tertiary age, the exact horizon being frequently uncertain. It may be convenient, however, to give a list of those woods which have been ascribed to earlier formations (excluding some doubtful records):

- Jurassic: *Suevioxylon*.
- Aptian: *Aptiana*, *Cantia*, *Dipterocarpoxyton porosum*, *Hythia*, *Sabulia*.
- Albian: *Laurinium albiense*.
- Cenomanian: *Hamamelidoxylon renaulti*, *Salicinium biradiatum*.
- Senonian: *Carpinoxylon compactum*, *Cornoxyton myricaeforme*, *C. vateri*, *Fegonium dryandraeforme*, *F. schenki*, *Juglandinium longiradiatum*, *Laurinium brunsvicense*, *Plataninium subaffine*, *Salicinium varians*, *Taenioxylon* sp.
- Upper Cretaceous: *Aachenoxylon*, *Caesalpinioxylon oweni*, *Dryoxylon americanum*, *Euonyminium auerbachi*, *Fegonium hokkaidense*, *Jugloxylon*, *Laurinium antiquum*, *L. haasi*, *L. uniseriatum*, *Nicolia aegyptiaca*, *N. moreneti*, *Nothofagoxylon scalariforme*, *Paraphyllanthoxylon*, *Populocaulis*, *Rhoidium ungeri*, *Rohlfisia*, *Sabiocaulis*, *Saururopsis*, *Sillimannia*.

The horizons given in the catalogue are usually as stated by the original authors, or as corrected by subsequent authors on the same subject. Many of them are doubtless capable of revision or of more accurate statement.

I have to thank Mr. F. M. Wonnacott and Dr. W. J. Jongmans for assistance in the preparation of this catalogue.

London, October, 1930.

[Published by permission of the Trustees
of the British Museum.]

Bibliography.

- Bailey, J. W. 1856. Letter from Professor J. W. Bailey, describing the Structure of the Fossil Plant from Posuncula River. U. S. Pacific Railroad Report, V, p. 337, pl. XII, figs. 1—2.
- Bailey, I. W. 1924. The problem of Identifying the Wood of Cretaceous and Later Dicotyledons: Paraphyllanthoxylon arizonense. Ann. Bot., XXXVIII, pp. 439—452, pl. XV.
- Barnard, C. 1927. A note on a Dicotyledonous Fossil Wood from Ulladulla, New South Wales. Proc. Linn. Soc. N. S. Wales, LII, II, pp. 113—121, pls. V, VI.
- Barron, T. 1905. On the Age of the Gebel Ahmar Sands and Sandstone, the Petrified Forest, and the Associated Lavas, between Cairo and Suez. Geol. Mag., (V), II, pp. 58—62.
- Bauer, J. 1927. Bestimmung der Stammpflanzen von Holzkohlen aus prähistorischen und subrezentem Fundorten Steiermarks. Oesterreich. Bot. Zeitschr., LXXXVI, pp. 29—42.
- Beck, R. 1882. Das Oligocän von Mittweida mit besonderer Berücksichtigung seiner Flora. Zeitschr. deutsch. Geol. Ges., XXXIV, pp. 735—770, pls. XXXI, XXXII.
- 1886. Beiträge zur Kenntnis der Flora des Sächsischen Oligocäns. Zeitschr. deutsch. Geol. Ges., XXXVIII, pp. 342—352, pl. VII.
- Becke, F. 1912. Fossiles Holz aus der Putzenwacke von Joachimstal. Min. u. Petr. Mitt., XXXI, pp. 81—86.
- Berger, L. G. den. 1923. Fossiele Houtsoorten uit het Tertiair van Zuid-Sumatra. Verhandl. Geol.-Mijnb. Gen. Nederland en Kol., VII, pp. 143—148.
- 1927. Unterscheidungsmerkmale von rezenten und fossilen Diptero-carpaceengattungen. Bull. Jard. Bot. Buitenzorg, (3), VIII, pp. 495—498.
- Berry, E. W. 1916. The Lower Eocene Floras of South-eastern North America. U. S. Geol. Surv. Prof. Paper 91, 481 pp., CXVII pls.
- 1918. The Fossil Higher Plants from the Canal Zone. Bull. U. S. Nat. Mus., 103, pp. 15—44, pls. XII—XVIII.
- 1922. Additions to the Flora of the Wilcox Group. U. S. Geol. Surv. Prof. Paper 131 A, pp. 1—21, pls. I—XVIII.
- 1924. The Middle and Upper Eocene Floras of South-eastern North America. U. S. Geol. Surv. Prof. Paper 92, 206 pp., LXIV pls.
- Blanckenhorn, M. 1901. Neues zur Geologie und Paläontologie Aegyptens. Zeitschr. deutsch. Geol. Ges., LIII, pp. 52—132.
- Bowerbank, J. S. 1844. On a New Variety of Vascular Tissue, found in a Fossil Wood from the London Clay. Trans. Micro. Soc. London, I, pp. 16—18, pl. II, figs. 1—5.
- Brongniart, A. 1822. Description des Végétaux fossiles du Terrain du Sédiment supérieur. In Cuvier & Brongniart, Descr. Géol. des Environs de Paris, ed. III, pp. 353—371.
- Carruthers, W. 1870. On the Petrified Forest near Cairo. Geol. Mag., (I), VII, pp. 306—310, pl. XIV.

- 1878. Description of Tertiary Plant-remains from Bracklesham and Worthing, Sussex. In Dixon and Jones, *Geology of Sussex*, ed. II, pp. 162—167, pl. XVI.
- Caspary, R. 1888. Einige fossile Hölzer Preussens. *Schriften Phys.-Oekonom. Ges. Königsberg*, XXVIII, pp. 27—45.
- 1889. [Edited by R. Triebel.] Einige fossile Hölzer Preussens. *K. Preuss. Geol. Landesanst., Abh. z. geol. Specialkarte von Preussen*, IX, 2, 86 pp., Atlas, XV pls.
- Chalk, L. & Rendle, B. J. 1929. *British Hardwoods: Their Structure and Identification*. *Forest Products Research Bull.*, 3, 53 pp., 45 figs.
- Chapman, F. 1918. On the Age of the Bairsdale Gravels; with a note on the included Fossil Wood. *Proc. Roy. Soc. Victoria*, (n. s.), XXXI, pp. 166—175, pl. X.
- Chhibber, H. L. 1928. The Origin and Mineral Constitution of the Late Tertiary Fossil Wood of Burma. *Journ. Proc. Asiatic Soc. Bengal*, XXIII, pp. 13—26, pls. I, II.
- Chiarugi, A. 1929. Legni Fossili. *Result. Sci. Miss. alle Oasi di Giarabub (1926—7)*, III, *Palaeontologia*, pp. 397—430, pls. XLV—XLVII.
- 1929a. Prime notizie sulle foreste petrificate della Sirtica. *N. Giorn. Bot. Ital.*, (n. s.), XXXV, pp. 558—566.
- Colani, M. 1919. Sur un Dipterocarpoxyylon annamense nov. sp. du Tertiaire supposé de l'Annam. *Bull. Serv. Geol. de l'Indochine*, VI, III, 8 pp., pls. I, II.
- Combes, P. 1907. Contribution à l'étude de la Flore Eocène. Sur un bois fossile nouveau appartenant à l'étage Sparnacien. *Bull. Soc. Géol. France*, (4), VII, pp. 28—29, pl. I.
- Compter, G. 1912. Fossile Hölzer aus dem Diluvium von Apolda. *Zeitschr. f. Naturwiss.*, LXXXIII, pp. 405—422.
- Conwentz, H. 1876. Ueber die Versteineten Hölzer aus dem Norddeutschen Diluvium. *Inaug. Diss.*, 34 pp., Breslau.
- 1880. Die Fossilen Hölzer von Karlsdorf am Zobten. *Schriften Nat. Ges. Danzig*, (N. F.), IV, 4, pp. 1—48, pls. I—VIII.
- 1882. Fossile Hölzer aus der Sammlung der Königlichen geologischen Landesanstalt zu Berlin. *Jahrb. Kgl. Geol. Landesanst.*, 1881, [Abh. 2], pp. 144—171.
- 1885. Sobre algunos árboles fósiles del Rio Negro. *Bol. Ac. Nac. Cienc. Cordoba*, VII, pp. 435—456.
- Corda, A. J. 1845. Beiträge zur Flora der Vorwelt. VIII + 128 pp., LX pls., Prague.
- Cramer, C. 1868. Fossile Hölzer der Arctischen Zone. *Flora Fossilis Arctica*, I, pp. 167—180, pls. XXXIV—XLII.
- Crié, L. 1888. Recherches sur la Flore Pliocène de Java. *Samml. Geol. Reichs. Mus. Leiden*, (1), V, pp. 1—22, pls. I—VIII.
- 1889. Beiträge zur Kenntnis der fossilen Flora einiger Inseln des Südpacifischen und Indischen Oceans. *Pal. Abh.*, V, 2, pp. 77—90, pls. IX—XVIII.
- Dawson, J. W. 1873. Note on the Fossil Plants from British Columbia collected by Mr. James Richardson in 1872. *Geol. Surv. Canada Progr. Rep.*, 1872—73, pp. 66—71. [Reprinted in *Amer. Journ. Sci.*, (3), VII, 1874, pp. 47—51.]
- 1875. Note on the Plants collected by Mr. G. M. Dawson from the Lignite Tertiary Deposits near the Forty-ninth Parallel. *Rep. Geol. & Resources 49th. Par.*, Appendix A, pp. 327—331.
- 1888. Note on Fossil Woods and other Plant Remains from the Cretaceous and Laramie Formations of the Western Territories of Canada. *Trans. Roy. Soc. Canada*, V, Sect. IV, pp. 31—37.
- Dixon, F. 1850. *The Geology and Fossils of the Tertiary and Cretaceous Formations of Sussex*. XVI+424+XVI pp., XL pls., London.

- Donath, M. 1928. Ein Fossiler Baumstamm als Einschluss im Basalt. *Centralbl. f. Min.*, A, 1928, pp. 57—61, 3 figs.
- Eames, A. J. 1910. On the Origin of the Broad Ray in *Quercus*. *Bot. Gaz.*, XLIX, pp. 161—167, pls. VIII, IX.
- Edwards, W. N. 1925. Report on Fossil Wood from Somaliland. *Mon. Geol. Dept. Hunterian Museum, Glasgow Univ.* I. The Collection of Fossils and Rocks from Somaliland, p. 171.
- Eichwald, E. 1865. *Végétaux Fossiles de la Période Moyenne. Le-thaea Rossica*, II, 1, pp. 1—71.
- Engelhardt, H. 1870. Flora der Braunkohlenformation im Königreich Sachsen. *Preisschr. Fürst. Jablonowsk. Ges. Leipzig*, XVI, 70 pp., atlas, XV pls.
- Ettingshausen, C. 1851. See Hauer.
- Falqui, G. 1906. Su Alcune Piante fossili della Sardegna. 25 pp., 1 pl., Cagliari.
- Felix, J. 1882. Studien über Fossile Hölzer. *Inaug. Diss.*, 84 pp., 1 pl., Leipzig.
- 1883. Untersuchungen über fossile Hölzer. *Zeitschr. deutsch. Geol. Ges.*, XXXV, pp. 59—91, pls. II, III.
- 1883a. Die fossilen Hölzer Westindiens. *Sammlung Pal. Abh.* I, pp. 1—29, V pls.
- 1884. Die Holzopale Ungarns in Palaeophytologischer Hinsicht. *Mitt. Jahrb. Ungar. Geol. Anstalt*, VII, pp. 1—43, IV pls.
- 1886. Untersuchungen über fossile Hölzer. II. *Zeitschr. Deutsch. Geol. Ges.* XXXVIII, pp. 483—492, pl. XII.
- 1887. Untersuchungen über fossile Hölzer. III. *Zeitschr. deutsch. Geol. Ges.*, XXXIX, pp. 517—523, pl. XXV.
- 1887a. Beiträge zur Kenntnis der Fossilen Hölzer Ungarns. *Mitt. Jahrb. Ungar. Geol. Anstalt*, VIII, 5, pp. 145—162, pls. XXVII, XXVIIa.
- 1893. Fossile Hölzer von Tlacolula. In Felix & Lenk, *Beiträge zur Geologie und Paläontologie der Republik Mexico*, II, 1, pp. 46—51, pl. III.
- 1894. Untersuchungen über fossile Hölzer. IV, (Hölzer aus dem Kaukasus). *Zeitschr. Deutsch. Geol. Ges.*, XLVI, pp. 79—110, pls. VIII—X.
- 1896. Untersuchungen über fossile Hölzer, V. *Zeitschr. deutsch. Geol. Ges.*, XLVIII, pp. 249—260, pl. VI.
- Fietz, A. 1926. Fossile Hölzer aus Schlesien. *Jahrb. Geol. Bundesanst.*, LXXVI, pp. 217—244, pl. V.
- 1926a. Prähistorische Holzkohlen aus der Umgebung Brünns. I Teil. *Planta: Archiv. f. Wiss. Bot.*, II, pp. 414—423.
- Fliche, P. 1888. Sur les bois silicifiés de la Tunisie et de l'Algérie. *C. R. Ac. Sci.*, CVII, pp. 569—572.
- 1898. Note sur les Bois Fossiles de Mételin. *Ann. des Mines*, [9], Méms. XIII, pp. 293—303.
- 1899. Note sur un bois de Vigne des cinérites du Cantal. *Bull. Soc. Géol. France*, [3], XXVII, pp. 318—321.
- 1905. Note sur des bois fossiles de Madagascar. *Bull. Soc. Géol. France*, [4], V, pp. 346—358, pl. X.
- Foxworthy, F. W. 1907. Philippine Woods. *Philippine Journ. Sci.*, C, Bot. II, pp. 351—404, 55 figs.
- 1909. Indo-Malayan Woods. *Philippine Journ. Sci.*, C, Bot. IV, pp. 409—592, pls. XXII—XXX.
- Fritel, P. H. 1909. Revision de la Flora Fossile des Grès Yprésiens du Bassin de Paris. *Journ. Bot.*, [2], II, pp. 149—169, 13 figs.
- & Viguier, R. 1911. Etude anatomique de deux bois éocènes. *Ann. Sci. Nat. Bot.*, [9], XIV, pp. 63—80, pl. I.
- Fujii, K. See Stopes & Fujii.

- Goeppert, H. R. 1839. Bemerkungen über die als Geschiebe im nördlichen Deutschland vorkommenden Versteinerten Hölzer. Neues Jahrb., pp. 518—521, pl. VIII B.
- 1845. See Goeppert & Berendt.
- 1854. Die Tertiärflora auf der Insel Java. 169 pp., XIV pls., 's Gravenhage.
- 1862. Ueber die in der Geschiebformation vorkommenden versteinerten Hölzer. Zeitschr. deutsch. Geol. Ges., XIV, pp. 551—554.
- 1881. Beiträge zur Pathologie und Morphologie fossiler Stämme. Palaeontographica, XXVIII, pp. 131—140, pls. I—V.
- & Berendt, G. C. 1845. Der Bernstein und die in ihm befindlichen Pflanzenreste der Vorwelt. 126 pp., VII pls., Berlin.
- Gothan, W. 1908. Die fossilen Hölzer von den Seymour und Snow-Hill Inseln. Wiss. Ergebn. Schwed. Südpol. Exped., 1901—03, III, VIII, 33 pp., II pls.
- 1910. Die Fossilen Holzreste von Spitzbergen. K. Svensk. Vet. Handl., XLV, 8, 56 pp., VII pls.
- Graves, L. 1847. Essai sur la Topographie Géognostique du Département de l'Oise. XV+804 pp., Beauvais.
- Hauer, F. 1851. Der Goldbergbau von Vöröspatak in Siebenbürgen. Jahrb. Geol. Reichs. Wien, II, 4, pp. 64—93.
- Heurn, J. F. C. van. 1927. Waarnemingen in verband met het voorkomen van verkieseld hout in de voormalige residentie Batavia. Handelingen Ned. Nat.-Gen. Congres Amsterdam, XXI, pp. 283—285, 1 pl.
- Hofmann, H. 1883. Ueber die fossilen Hölzer aus dem mecklenburgischen Diluvium. Arch. Ver. Fr. Nat. Mecklenburg, XXXVI, pp. 65—107.
- 1884. Untersuchungen über fossile Hölzer. Zeitschr. f. Naturwiss., LVII, pp. 156—195 [Also published separately, Inaug. Diss., Halle, 44 pp.]
- 1884a. Verkieselte Hölzer aus Aegypten. Zeitschr. f. Naturwiss., LVII, pp. 484—486, pl. III, figs. 1, 2.
- Hofmann, E. 1928. Verkieselte Hölzer aus dem Museum in Szombathely. Ann. Comit. Castriferrei (Hist. Nat.), A, 1928. (pp. 1—4, 1 pl.).
- 1929. Verkieselte Hölzer von der Vashegy (Eisenberg)-Gruppe. Ann. Sabarienses, III, pp. 81—87, pls. I—IV.
- Holden, R. 1916. A Fossil Wood from Burma. Rec. Geol. Surv. India, XLVII, pp. 267—272, pl. XXIX.
- Hopkinson, A. D. 1912. Beiträge zur Mikrographie tropischer Hölzer. Beihefte Bot. Centralbl., XXIX, 2, pp. 441—456, 24 text-figs.
- Houlbert, C. 1893. Recherches sur la Structure Comparée du Bois Secondaire dans les Apétales. Ann. Sci. Nat. Bot., (7), XVII, pp. 1—184, pls. I—VIII.
- 1910. Les Bois des Faluns de Touraine. Feuille Jeunes Nat., XL, pp. 69—76, pls. III—VIII.
- Hovelacque, M. 1889. Sur la nature Végétale de l'Aachenosaurus. (résumé). Bull. Soc. Belge Géol., III, pp. 505, 506.
- 1890. Sur la Nature Végétale de l'Aachenosaurus multident G. Smets. Bull. Soc. Belge Géol., IV, pp. 59—72, pl. III, 8 text-figs.
- Janssonius, H. H. See Moll & Janssonius.
- Jeffrey, E. C. & Chrysler, M. A. 1906. The Lignites of Brandon. Rept. Vermont State Geol., 1905—6, pp. 195—201, pls. XLIX—LI.
- Johnson, M. H. 1876. On Silicified Structure in Pyritized Wood. Journ. Quekett Micro. Club, IV, pp. 159—160.
- Johnston, R. M. 1875. The Launceston Tertiary Basin. Papers & Proc. Roy. Soc. Tasmania, 1874, pp. 53—62, 1 pl.
- 1888. Systematic Account of the Geology of Tasmania. XXII+408 pp., LVII pls., Hobart.

- K a i s e r, P. 1879. Ulmoxyton. Ein Beitrag zur Kenntnis fossiler Laubhölzer. Zeitschr. f. Gesamt. Naturwiss., [3], IV, pp. 88—100.
- 1880. Ficoxyton bohemicum, ein neues fossiles Laubholz. Zeitschr. f. Gesamt. Naturwiss., [3], V, pp. 309—317.
- 1880a. Neue fossile Laubhölzer. Bot. Centralbl., pp. 511—512.
- 1890. Die fossilen Laubhölzer. Wiss. Beilage Jahresber. Realprogymn. Schönebeck a. E., 46 pp.
- K a n e h i r a, R. 1921. Anatomical Characters and Identification of Formosan Woods. 317 pp., L pls., Taihoku.
- 1921a. Identification of the Important Japanese Woods by Anatomical Characters. 104 pp., IX pls., Taihoku.
- K n o w l t o n, F. H. 1891. Description of Fossil Woods and Lignites from Arkansas. Arkansas Geol. Surv. Ann. Rept., 1889, 2, pp. 249—268, pls. IX—XI.
- 1899. Fossil Flora of the Yellowstone National Park. Mon. U. S. Geol. Surv., XXXII, II, pp. 651—882, pls. LXXVII—CXXI.
- 1902. Notes on the Fossil Fruits and Lignites of Brandon, Vermont. Bull. Torrey Bot. Club, XXIX, pp. 635—641, pl. XXV.
- K o b b e, F. 1887. Ueber fossile Hölzer der mecklenburger Braunkohle. Arch. Ver. Fr. Nat. Mecklenburg, XLI, pp. 89—142, pls. II—III.
- K r a u s, G. 1865. Ueber einige bayerische Tertiärhölzer. Würzburger Naturwiss. Zeitschr., VI, pp. 45—48.
- 1882. Ueber ein tertiäres Laubholz aus den sicilianischen Schwefelgruben. Ber. Sitz. Nat. Ges. Halle, pp. 4—5.
- 1882a. Ueber das Holz des mexikanischen Chijol. Ibid., pp. 5—7.
- 1882b. Ueber das Sandelholz von Juan Fernandez. Ibid., pp. 8—9.
- 1882c. Beiträge zur Kenntnis der fossilen Hölzer. I. Hölzer aus den Schwefelgruben Siciliens. Abh. Nat. Ges. Halle, XVI, pp. 77—91, pl. I.
- K r ä u s e l, R. 1920. Nachträge zur Tertiärflora Schlesiens. II. Braunkohlenhölzer. Jahrb. Preuss. Geol. Landesanst., XXXIX, 1, f. 1918, pp. 418—460, pls. XXVIII—XXXVIII.
- 1922. Fossile Hölzer aus dem Tertiär von Süd-Sumatra. Verh. Geol. Mijnb. Gen. Nederland en Kol., V, pp. 231—287, VII pls.
- 1922a. Ueber einen fossilen Baumstamm von Bolang (Java). Proc. Sect. Sci. XXV, pp. 9—15, pl. I. [From Versl. Gew. Vergad. Wis. Nat. Afd., Akad. v. Wetensch. Amsterdam, XXXI.]
- 1924. Beiträge zur Kenntnis der fossilen Flora Südamerikas. I. Fossile Hölzer aus Patagonien und benachbarten Gebieten. Arkiv f. Bot., XIX, 9, 36 pp., IV pls.
- 1925. Der Stand unserer Kenntnisse von der Tertiärflora Niederländisch-Indiens. Verhandl. Geol. Mijnb. Gen. Nederland en Kol., VIII, pp. 329—342.
- 1926. Ueber einige Fossile Hölzer aus Java. Leidsche Geol. Med., II, 1, pp. 1—6, pls. I, II.
- 1928. Ueber ein Juraholz vom Angiospermentypus. Senckenbergiana, X, pp. 250—254, figs. 5—8.
- K r e n d o w s k y, M. 1880. Beschreibung der fossilen Bäume hauptsächlich aus dem Süden Russlands. [In Russian]. Arb. Nat. Ges. Univ. Kharkov, XIII. [Not seen].
- K u n t z e, O. 1895. Verkieselungen und Versteinerungen von Hölzern. Geogenetische Beiträge, Leipzig, pp. 24—36.
- L a k o w i t z, 1890. Betuloxyton geinitzii nov. sp. und die fossilen Birkenhölzer. Schriften Nat. Ges. Danzig, (N. F.), VII, 3, pp. 25—32, I pl.
- L a n g e, T. 1890. Beiträge zur Kenntnis der Flora des Aachener Sandes. Zeitschr. deutsch. Geol. Ges., XLII, pp. 658—676, pls. XXXI—XXXIV.

- Lignier, O. 1907. Végétaux Fossiles de Normandie IV. Bois Divers (1er serie). Mém. Soc. Linn. Normandie, XXII, pp. 237—332, pls. XVII—XXIII.
- Lingelsheim, A. 1917. Ein Beitrag zur fossilen Flora Ungarns. Jahrb. Kgl. Ungar. Geol. Reichsanst., 1915, pp. 545—563, 13 text-figs.
- Massalongo, A. 1852. Sapindacearum Fossilium Monographia. 28 pp., VI pls., Verona.
- Meneghini, J. 1857. Paléontologie de l'Île de Sardaigne. 586 pp., VIII pls., Turin.
- Mercklin, C. 1855. Palaeodendrologikon Rossicum. 99 pp., XX pls. St. Petersburg.
- Meschinelli, L. & Squinabol, X. 1893. Flora Tertiaria Italica. 578 pp., Padova.
- Moeller, J. 1876. Beiträge zur vergleichenden Anatomie des Holzes. Denkschr. Ak. Wiss. Wien, XXXVI, pp. 297—426, pls. I—VI.
- Moll, J. W. & Janssonius, H. H. 1906. Mikrographie des Holzes der auf Java vorkommenden Baumarten. Leiden. (Lief. 1—IX, 1906—1928. The work is not yet complete.)
- Moll, J. W. & Janssonius, H. H. 1912. The Linnean Method of Describing Anatomical Structures. Some Remarks concerning the Paper of Mrs. Dr. Marie C. Stopes, entitled „Petrifications of the Earliest European Angiosperms.” Proc. Sect. Sci. XV, 1, pp. 620—629, 2 textfigs. [From Versl. Gew. Vergad. Wis. Nat. Afd., Akad. v. Wetensch. Amsterdam, XXI.]
- Muller, N. J. C. 1888. Atlas der Holzstruktur dargestellt in Microphotographien. Mit erläuterndem Text. 110 pp., 63 figs., Atlas XXI pls.
- Nagelhard, K. 1922. (Formerly Nagel). Ulmaceae. Foss. Cat. Plantae, pars 10, 84 pp.
- Nagel, K. 1915. Juglandaceae. Foss. Cat. Plantae, pars 6, 87 pp. — 1916. Betulaceae. Foss. Cat. Plantae, pars 8, 177 pp.
- Nicol, W. 1835. On the Structure of some Fossil Woods found in the Island of Mull, Northern Africa, and on the Karoo Ground to the North-East of the Cape of Good Hope. Edinburgh New Phil. Journ., XVIII, pp. 335—338.
- Oswald, F. 1914. The Miocene Beds of the Victoria Nyanza. Quart. Journ. Geol. Soc., LXX, pp. 128—162.
- Pampaloni, L. 1902. Sopra alcuni tronchi Silicizzati dell' Eocene superiore dell' Impruneta (Provincia di Firenze). Boll. Soc. Geol. Ital., XXI, pp. 25—29, pl. I.
- 1904. Sopra alcuni Legni Silicizzati del Piemonte. Boll. Soc. Geol. Ital., XXII, pp. 535—548, text-figs. 1—14.
- Pax, F. 1901. Aceraceae. Engler's Pflanzenreich. 89 pp.
- Penhallow, D. P. 1903. Notes on Tertiary Plants. Trans. Roy. Soc. Canada, (2), IX, Sect. IV, pp. 33—96, 29 figs.
- 1907. Notes on Fossil Woods from Texas. Trans. Roy. Soc. Canada, (3), I, Sect. IV, pp. 93—114, 8 figs.
- 1907a. A Report on Fossil Plants from the International Boundary Survey for 1903—05. Trans. Roy. Soc. Canada, (3), I, Sect. IV, pp. 287—334, pls. I—IX.
- 1907b. Contributions to the Pleistocene Flora of Canada. Amer. Nat., XLI, pp. 443—452.
- 1908. Report on Tertiary Plants of British Columbia. Canada Dept. Mines, Geol. Survey Branch. No. 1013, 167 pp.
- Pfeiffer, J. P. & Heurn, J. F. C. van. 1928. Some Fossil Woods from Java not yet described. Proc. Sect. Sci., XXXI, 2, pp. 1005—1011, II pls. [From Versl. Gew. Vergad. Wis. Nat. Afd., Akad. v. Wetensch. Amsterdam, XXXVII].

- Platen, P. 1908. Untersuchungen fossiler Hölzer aus dem Westen der Vereinigten Staaten von Nordamerika. XVI+155 pp., III pls., Leipzig.
- Prokesch-Osten & Unger, F. 1852. Die versteinerten Holzstämmen im Hafen von Sigri auf der Insel Lesbos. Sitzber. Ak. Wiss. Wien, IX, pp. 855—858.
- Quenstedt, A. 1885. Handbuch der Petrefaktenkunde, 3. Aufl. Tübingen, 1240 pp., atlas, C pls.
- Reiss, K. 1907. Untersuchung über fossile Hölzer aus Japan. 224 pp., 1 pl., Inaug. Diss. Rostock.
- Rendle, B. J. See Chalk, L.
- Roemer, F. 1852. See Unger.
- Sahni, B. 1920. Petrified Plant Remains from the Queensland Mesozoic and Tertiary Formations. Queensland Geol. Survey, no. 267, 48 pp., V pls.
- Schaeffer, G. C. 1856. Description of the Structure of the Fossil Wood from the Colorado Desert. U. S. Pacific Railroad Rep., V, pp. 338—339, pl. XII, figs. 3—4.
- Schenk, A. 1869. Ueber einige in der Braunkohle Sachsens vorkommende Pflanzenreste. Bot. Zeitung, XXVII, pp. 375—380.
- 1880. Ueber fossile Hölzer aus der Libyschen Wüste. Bot. Zeitung, XXXVIII, pp. 657—661.
- 1883. Fossile Hölzer. (Beiträge zur Geol. u. Pal. der Libyschen Wüste). Palaeontographica, XXX, 2, Abt. I, pp. 1—19, pls. 1—V.
- 1888. Fossile Hölzer aus Ostasien u. Aegypten. Bihang K. Svensk. Vet. Akad. Handl., XIV, Afd. III, no. 2, 24 pp.
- 1890. Dicotyle Hölzer. Palaeophytologie, II, 9, pp. 890—904.
- Schimper, W. P. 1872. Traité de Paléontologie Végétale, II. 968 pp., Paris.
- 1874. Ibid., III. 896 pp., Paris.
- Schleiden see Schmid & Schleiden.
- Schmid, E. 1853. Die organischen Reste des Muschelkalkes im Saal-Thale bei Jena. Neues Jahrb., 1853, pp. 9—30.
- Schmid, E. E. & Schleiden, M. J. 1855. Ueber die Natur der Kieselhölzer. 42 pp., III pls., Jena.
- Schönfeld, G. 1919. Ueber einige neue Funde fossiler Hölzer aus der Umgebung Dresdens. Abh. Nat. Ges. Isis, Dresden, 1918, pp. 3—16, pl. I.
- 1925. Ueber unsere Braunkohlenwälder und die Entstehung der Braunkohlenflöze. Ber. Freiburger Geol. Ges., X, pp. 18—24.
- Schönfeld, E. 1930. Ueber zwei neue Laubhölzer aus dem Miozän. Senckenbergiana, XII, pp. 111—126, figs. 1—18.
- 1930a. Sächsische Braunkohlenhölzer. Untersucht und bestimmt von G. Schönfeld. Sitzber. Nat. Ges. Isis, Dresden, 1929, pp. 68—70.
- Schroeter, C. 1880. Untersuchung über Fossile Hölzer aus der arctischen Zone. Flora Fossilis Arctica, VI, 1, 38 pp., III pls.
- Schuster, J. 1908. Ueber ein fossiles Holz aus dem Flysch des Tegernseer Gebietes. Geognost. Jahreshfte, XIX, pp. 139—152, pl. II, 3 text-figs.
- 1908a. Ueber ein Pliocänes Eichenholz aus Idaho. Neues Jahrb. 1908, II, pp. 49—54, pls. III, IV.
- 1909. Paläobotanische Notizen aus Bayern. 5. Ueber ein oberoligozänes Lorbeerholz aus dem Algäu. Ber. Bayer. Bot. Ges., XII, I, pp. 54—56.
- 1910. Ueber Nicolien und Nicolien ähnliche Hölzer. K. Svensk. Vet. Akad. Handl., XLV, 6, 18 pp., III pls.
- 1911. Paleozäne Rebe von der Greifswalder Oie. Ber. deutsch. Bot. Ges., XXIX, pp. 540—544, pl. XX.

- 1911a. Die Flora der Trinil-Schichten. Die Pithecanthropus-Schichten auf Java. Leipzig, pp. 235—257, pls. XXXI, XXXII, 8 text-figs.
- Schweinfurth, G. 1882. Zur Beleuchtung der Frage über den versteinerten Wald. Zeitschr. deutsch. Geol. Ges., XXXIV, pp. 139—145.
- Seward, A. C. & Holttum, R. E. 1924. Tertiary Plants from Mull. In E. B. Bailey & others, Tertiary and Post-Tertiary Geology of Mull, Loch Aline, and Oban. Mem. Geol. Surv. Scotland, pp. 67—90, pl II, 15 figs.
- Smets, G. 1888. *Aachenosaurus multidentis*, Reptile fossile des Sables d'Aix-la-Chapelle. Mém. Congr. Sav. Cathol. Paris. [Not seen].
- Solereder, H. 1899. Systematische Anatomie der Dicotyledonen. 984 pp., Stuttgart. (Ergänzungsband, 1908, 422 pp.)
- 1908. Systematic Anatomy of the Dicotyledons. (Transl. Boodle & Fritsch). 2 vols., 1182 pp., Oxford.
- Sorby, H. C. 1852. On the Occurrence of Non-Gymnospermous Exogenous Wood in the Lias, near Bristol. Trans. Mier. Soc. London, III, pp. 91—92, pl. XVI.
- Stache, G. 1855. De Casuarinis nunc viventibus et fossilibus nonnulla. 42 pp. Inaug. Diss., Vratislaviae.
- St. John, R. N. 1927. Replacement vs. Impregnation in Petrified Wood. Econ. Geol., XXII, pp. 729—739.
- Staub, M. 1889. Magyarorszag kövesült fatörzsei. (On Hungarian fossil woods.) Potf. Termesz. Közl. Budapest, VIII, pp. 182—191.
- Stokes, C. 1837. Notice Respecting a Piece of Recent Wood partly petrified by Carbonate of Lime, with some remarks on Fossil Woods. Trans. Geol. Soc. London, (2), V, pp. 207—214, pls. XVI, XVII.
- Stone, H. 1904. The Timbers of Commerce and their Identification. 311 pp., 183 figs., London.
- 1921. A Text-Book of Wood. 240 pp., XLI pls., London.
- Stopes, M. C. 1912. Petrifications of the Earliest European Angiosperms. Phil. Trans. Roy. Soc., London, B, CCIII, pp. 75—100, pls. VI—VIII.
- 1915. The Cretaceous Flora, Pt. II, Lower Greensand (Aptian) Plants of Britain. Cat. Mesozoic Plants, Brit. Mus. (Nat. Hist.), 360 pp., XXXII pls.
- Stopes, M. C. & Fujii, K. 1910. Studies on the Structure and Affinities of Cretaceous Plants. Phil. Trans. Roy. Soc., London, B, CCI, pp. 1—90, pls. I—IX.
- Stromer, E. 1924. Die Pflanzen-führenden Schichten Aegyptens. Abh. Bayer. Ak. Wiss., Math.-Nat. Abt., XXX, 2, pp. 5—27.
- Stur, D. 1873. Beiträge zur genaueren Deutung der Pflanzenreste aus dem Salzstocke von Wieliczka. Verh. Geol. Reichsanst. Wien, pp. 6—10.
- Szafer, W. 1914. Anatomische Bestimmung der diluvialen Holzreste von Ludwinow. Bull. Int. Acad. Sci. Cracovie, 2 B, pp. 345—350.
- Tchihatchef, P. 1863. Note sur deux nouveaux genres de bois fossile recueillis dans les environs de Constantinople. C. R. Ac. Sci., Paris, LVI, pp. 516—518.
- Tetzlaff, G. 1881. Die Holzreste von Leubingen. Ein Beitrag zur Kenntnis der deutschen Laubhölzer mit besonderer Berücksichtigung der fossilen. 44 pp., Inaug. Diss., Halle.
- Thomas, P. 1893. Description de quelques Fossiles Nouveaux ou Critiques des Terrains Tertiaires et Secondaires de la Tunisie. Végétaux Fossiles, pp. 1—4 [Mainly reprinted from Fliche 1888]. Expl. Sci. de la Tunisie. Paris.

- Triebel, R. 1889. See Caspary (Posthumous work, edited by Triebel).
- Tuzson, J. 1909. Anatomische Bestimmung der fossilen Hölzer des Balatonsees. *Math. Nat. Ber. Ungarn*, XXIV, p. 376.
- 1911. Monographie der Fossilen Pflanzenreste der Balatonseegegend. Resultate der wiss. Erforsch. des Balatonsees, I, I, Pal. Anhang IV, I, 64 pp., II pls.
- Udluft, H. 1926. Zeolithe als Fossilisationsmaterial. *Arkiv f. Kemi, Min. och Geol.*, IX, no. 33, 15 pp., II pls.
- Unger, F. 1842. Synopsis Lignorum Fossilium Plantarum Acramphibryarum. In Endlicher's *Genera Plantarum Suppl. II*, Appendix, pp. 100—102.
- 1842a. Ueber die Untersuchung fossiler Stämme holzartiger Gewächse. *Neues Jahrb.*, 1842, pp. 149—178. [Pp. 172—178 appear to be a reprint of Unger, 1842.]
- 1842b. Ueber die versteinerten Hölzer des National-Museums zu Linz. *Neues Jahrb.*, 1842, pp. 745—748.
- 1845. Synopsis Plantarum fossilium. XVIII + 330 pp. Leipzig. [Pp. 1—330 are identical with pp. XXV—CX of Unger's *Chloris Protogaea*.]
- 1847. *Chloris Protogaea*. CX + 150 pp., L pls., Leipzig, (1841—1847).
- 1849. Die Pflanzenreste im Salzstocke von Wieliczka. *Denkschr. Ak. Wiss. Wien*, I, pp. 311—322, pl. XXXV.
- 1850. *Genera et Species Plantarum Fossilium*. 628 pp., Vindobonae.
- 1852. *Iconographia Plantarum Fossilium*. *Denkschr. Ak. Wiss. Wien*, IV, pp. 73—118, pls. XXIV—XLV.
- 1852a. Beschreibung fossiler Hölzer aus Texas. In Roemer, F. *Die Kreidebildungen von Texas und ihre organischen Einschlüsse*, pp. 94—95, Bonn.
- 1852b. See Prokesch-Osten.
- 1854. Die Fossile Flora von Gleichenberg. *Denkschr. Ak. Wiss. Wien*, VII, pp. 157—184, pls. I—VIII.
- 1857. Ueber fossile Pflanzen des Süßwasserkalkes und Quarzes. *Denkschr. Ak. Wiss. Wien*, XIV, pp. 1—12, pls. I—III.
- 1857a. Beiträge zur näheren Kenntnis des Leithakalkes. *Denkschr. Ak. Wiss. Wien*, XIV, pp. 13—38, pls. IV, V.
- 1859. Der versteinerte Wald bei Cairo und einige andere Lager verkieselten Holzes in Aegypten. *Sitzber. Ak. Wiss. Wien*, XXXIII, pp. 209—232, pls. I—III.
- 1863. See Tchihatchef.
- 1864. Fossile Pflanzenreste aus Neu-Seeland. In Hochstetter, *Novara Exped.*, *Geol. I*, 2, pp. 1—13, pls. I—V.
- 1866. Notiz über fossile Hölzer aus Abyssinien. *Sitzber. Ak. Wiss. Wien*, LIV, 1, pp. 289—297, pl. I.
- 1866a. Flore Tertiaire Moyenne. In Tchihatchef, *Asie Mineure, Paléontologie*, pp. 319—325, pl. XVII.
- Vater, H. 1884. Die Fossilen Hölzer der Phosphoritlager des Herzogthums Braunschweig. *Zeitschr. deutsch. Geol. Ges.*, XXXVI, pp. 783—853, pls. XXVII—XXIX.
- Viguier, see Fritel.
- Warren, E. 1912. On some specimens of Fossil Woods in the Natal Museum. *Ann. Natal Museum*, II, pt. III, p. 345—380, pls. XXV—XXVII.
- Watelet, A. 1866. *Description des Plantes Fossiles du Bassin de Paris*, 264 pp., Atlas LX pls., Paris.
- Wetzel, W. 1913. Ueber ein Kieselholzgeschiebe mit Teredonen aus den Holtenauer Kanal-Aufschlüssen. *Jahresber. Niedersächs. Geol. Ver.*, VI, pp. 20—59, pls. 1—III.

- Wilhelm, K. 1903. Laubhölzer. In Wiesner, Die Rohstoffe des Pflanzenreiches, 2. Aufl., II, pp. 872—1020.
- Windisch, P. 1886. Beiträge zur Kenntnis der Tertiärflora von Island. 52 pp. Inaug. Diss., Halle.
- Witham, H. T. M. 1833. The Internal Structure of Fossil Vegetables. 84 pp., XVI pls., Edinburgh.
- Wittmack, L. & Buchwald, J. 1902. Pflanzenreste aus der Hünenburg bei Rinteln a. d. Weser, und eine verbesserte Methode zur Herstellung von Schnitten durch verkohlte Hölzer. Ber. deutsch. Bot. Ges., XX, pp. 21—31, pl. III.
- Zander, R. 1923. Ein Beitrag zur Kenntnis der Tertiären Braunkohlenhölzer des Geiseltals. Braunkohle XXII, pp. 17—25, 38—42, pls. I—III.
-

Aachenoxylon Hovelacque, 1890, p. 60.

Aachenoxylon multidentis (Smets).

1888 *Aachenosaurus multidentis* Smets, pars. [Not seen].

1890 *Aachenoxylon* sp.: Hovelacque, p. 60, pl. III, f. 1; text-fig. 1.

Remarks: Originally described as the dermal spine of a reptile, but shown to be a dicotyledon by its internal structure. Affinities uncertain. Another supposed part of "*Aachenosaurus*" was described by Hovelacque as *Nicolia moresneti*.

Occurrence: Upper Cretaceous; Belgium (Moresnet).

Acacioxylon Schenk, 1883, p. 9.

Acacioxylon antiquum Schenk.

1883 *Acacioxylon antiquum* Schenk, p. 9.

1888 *Acacioxylon antiquum* Schenk: Fliche, p. 571.

1890 *Acacioxylon antiquum* Schenk: Kaiser, p. 31.

1893 *Acacioxylon antiquum* Schenk: Thomas, p. 3.

Occurrence: Lower Oligocene (Gebel Ahmar beds); Libyan Desert (Wadi Dugla).

Tertiary; Tunisia (Ain Cherichira).

Acacioxylon tenax Felix.

1893 *Acacioxylon tenax* Felix, p. 49, pl. III, t. 4, 6-8.

Occurrence: ? Tertiary; Mexico.

Acacioxylon vegae Schenk.

1888 *Acacioxylon vegae* Schenk, p. 23.

Remarks: Closely related to *A. antiquum* Schenk.

Occurrence: Lower Oligocene?; Egypt.

Acer L.

(See also **Acerinium**.)

Acer sp.

1914 *Acer* sp.: Szafer, p. 349.

1926a *Acer*: Fietz, p. 419.

Occurrence: Pleistocene; Poland.

Prehistoric (Loess); Czechoslovakia (Brno).

***Acerinium* Unger, 1842, p. 101.**

***Acerinium aegyptiacum* Schenk.**

1888 *Acerinium aegyptiacum* Schenk, p. 22.

1890 *Acerinium aegyptiacum* Schenk: Schenk, p. 900, f. 431.

Occurrence: Tertiary; Egypt (eastern desert, between Cairo and Suez).

***Acerinium astianum* Pampaloni.**

1904 *Acerinium astianum* Pampaloni, p. 544, f. 7—9.

Occurrence: Tertiary; Piedmont (Astigiano alto).

***Acerinium borussicum* (Caspary).**

1888 *Acer borussicum* Caspary, p. 38.

1889 *Acer borussicum* Caspary: Caspary, p. 7, pl. II, f. 1—6.

1890 *Acer borussicum* Caspary: Kaiser, p. 27.

Remarks: Original in Mus. Bot. Gart., Königsberg. Pax (1901, p. 77) gives reasons for doubting whether this wood belongs to *Acer*.

Occurrence: ? Tertiary; East Prussia.

***Acerinium danubiale* Unger.**

1842 *Acerinium danubiale* Unger, p. 101.

1842a *Acerinium danubiale* Unger: Unger, p. 175.

1842b *Acerinium „danubiense”* Unger: Unger, p. 748.

1845 *Acerinium danubiale* Unger: Unger, p. 237.

1847 *Acerinium danubiale* Unger: Unger, p. 136, pl. XLIV, f. 9—11.

1850 *Acerinium danubiale* Unger: Unger, p. 453.

1890 *Acerinium danubiale* Unger: Kaiser, p. 27.

Occurrence: Tertiary; Upper Austria.

***Acerinium terrae-coeruleae* (Caspary).**

1888 *Acer terrae-coeruleae* Caspary, p. 38.

1889 *Acer terrae-coeruleae* Caspary: Caspary, p. 11, pl. II, f. 7, 8; III, f. 1—7; IV, f. 1—12.

1890 *Acer terrae-coeruleae* Caspary: Kaiser, p. 27.

Remarks: Original in Mus. Bot. Gart., Königsberg. Pax (1901, p. 77) gives reasons for doubting whether this wood belongs to *Acer*.

Occurrence: Oligocene; Prussia (Palmnicken).

***Alnoxylon* Felix, 1884, p. 10.**

(Including ***Rhizoalnoxylon*** Conwentz).

***Alnoxylon inclusum* (Conwentz).**

1880 *Rhizoalnoxylon inclusum* Conwentz, p. 38, pl. VII, f. 25—28.

1890 *Rhizoalnoxylon inclusum* Conwentz: Kaiser, p. 9.

Remarks: The use of the prefix *Rhizo-* for root woods, introduced by Conwentz for this and other genera (chiefly conifers), only increases nomenclatorial confusion and has not been generally adopted. Felix (1884, p. 11) points out the difficulties of this usage. These roots were found in a cupressineous rootstock.

Occurrence: Tertiary; Silesia (Karlsdorf).

Fossilium Catalogus II. 17.

Alnoxyton vasculosum Felix.1884 *Alnoxyton vasculosum* Felix, p. 10, pl. I, f. 1.1889 *Alnoxyton vasculosum* Felix: Staub, p. 187.1890 *Alnoxyton vasculosum* Felix: Kaiser, p. 10.Remarks: Originals in Mus. Ungar. Geol. Reichsanst. Felix rightly substitutes *Alnoxyton* for *Rhizoalnoxyton* Conwentz.

Occurrence: Tertiary (Pannonischer Schichten); Hungary (Gyepüfüzes).

Alnoxyton yezoëense Reiss.1907 *Alnoxyton yezoëense* Reiss, p. 194, pl. I, f. 9, 10.

Occurrence: Miocene; Japan (Ishikari).

Alnoxyton sp.1902 *Alnus* sp.: Pampaloni, p. 29, pl. I, f. 5.Remarks: Considered by Pampaloni to be an example of the *Betulaceae*, and probably of the genus *Alnus*.

Occurrence: Eocene; Italy (Impruneta nr. Florence).

Alnus L.**Alnus** sp.1882 *Alnus* sp.: Conwentz, p. 146.1914 *Alnus* sp.: Szafer, p. 349.

Occurrence: Pleistocene; Holstein (Büsum), Poland.

[**Ambaroxylon** Houlbert, 1910, p. 74].Synonym of **Liquidambaroxylon**, q. v.**Ampeloxylon** Fliche, 1899, p. 321.**Ampeloxylon ampelopsoides** (Schönfeld).1930 *Vitoxylon ampelopsoides* Schönfeld, p. 112, f. 1—9.Remarks: According to the author, this wood agrees most closely with the recent genus *Ampelopsis* among the *Vitaceae*. Fliche's form-genus *Ampeloxylon* has priority over *Vitoxylon* Schuster.

Occurrence: Miocene; "Deckgebirge der miozänen Braunkohle des Lettengrabens in der hohen Rhön."

Ampeloxylon cineritarum Fliche.1899 *Ampeloxylon cineritarum* Fliche, p. 321.Remarks: Leaves of *Vitis* have been recorded by Saporta from the same locality.

Occurrence: Pliocene; Cantal.

Ampeloxylon coheni (Schuster).1911 *Vitoxylon coheni* Schuster, p. 541, pl. XX.

Remarks: Fliche's generic name has priority.

Occurrence: Lower Eocene; Greifswalder Oie near Rügen.

Anacardioxylon Felix, 1882, p. 70.

Anacardioxylon magniporosum Platen.

1908 *Anacardioxylon magniporosum* Platen, p. 57.

Occurrence: Tertiary; California.

[*Anacardioxylon mollii* Kräusel.]

Synonym of *Sumatroxylon mollii*, q. v.

Anacardioxylon spondiaeforme Felix.

1882 *Anacardioxylon spondiaeforme* Felix, p. 70.

1883a *Anacardioxylon spondiaeforme* Felix: Felix, p. 16, pl. II, f. 7, 9.

1890 *Anacardioxylon spondiaeforme* Felix: Kaiser, p. 25.

Remarks: Similar to *Spondias lutea*. Originals in Munich.

Occurrence: Tertiary; Antigua.

Anacardioxylon uniradiatum Felix.

1894 *Anacardioxylon uniradiatum* Felix, p. 92, pl. VIII, f. 1a—e.

Remarks: Compared with *Spondias lutea*, but also with some *Simarubaceae*.

Occurrence: Eocene (Sumgait series); Caucasus (Apscheron).

Anomaloxylon Felix, 1887, p. 527 (non Gothan, 1910).

Remarks: Quite apart from the question of priority, Gothan's coniferous genus *Anomaloxylon* (Gothan, 1910, p. 11) cannot be regarded as well founded. The abnormal condition of the wood rays, which was the principal character of the genus, might be due to disease or wounding.

Gothan's species *A. magnoradiatum* was renamed *Protocedroxylon magnoradiatum* by Eckhold (Jahrb. Pr. Geol. Land., XLII, p. 491), which can hardly be correct if, as Gothan says, the ray cells do not show abietinean pitting. On another page of the same paper (p. 501) Eckhold puts the species, with a query, in his genus *Protocupressinoxylon*, and on p. 493 he gives both names. For the present, the latter might be used.

Anomaloxylon vicentinum Felix.

1887 *Anomaloxylon vicentinum* Felix, p. 527, pl. XXV, f. 8.

1890 *Anomaloxylon vicentinum* Felix: Kaiser, p. 38.

Remarks: A liane which, on grounds of certain anomalous structure of the wood, Felix separated from his form-genus *Helictoxylon*. Originals in Mus. Univ. Breslau.

Occurrence: Tertiary; Italy (Monte Grumi near Castelgomberto).

Aptiana Stopes, 1912, p. 84.

Aptiana radiata Stopes.

1912 *Aptiana radiata* Stopes, p. 84, pl. VI, f. 1, 3—5; VII, f. 6; VIII, f. 10, 11; text-fig. 1—5.

1912 *Aptiana radiata* Stopes: Moll & Janssonius, p. 622.

1915 *Aptiana radiata* Stopes: Stopes, p. 284, text-fig. 87—92.

1924 *Aptiana radiata* Stopes: Bailey, p. 448.

Remarks: Compared by Stopes with various members of the *Caprifoliaceae*, *Magnoliaceae*, *Aquifoliaceae* and *Cyrillaceae*. Moll & Janssonius concluded that it belonged to the *Ternstroemiaceae*, and was very close to the living *Eurya acuminata*. Stopes (1915) did not entirely agree with this. Bailey suggested a comparison with *Vaccinium*. Originals in Geol. Dept., Brit. Mus. (Nat. Hist.). Also figured Scott, D. H. 1924. *Extinct Plants*, p. 55, f. 7.

Occurrence: Lower Cretaceous (Aptian); England (Isle of Wight).

Aralinium Platen, 1908, p. 59.

Aralinium excellens Platen.

1908 *Aralinium excellens* Platen, p. 59.

Occurrence: Mio-Pliocene; California.

Aralinium lindgreni Platen.

1908 *Aralinium lindgreni* Platen, p. 64.

Occurrence: Mio-Pliocene; California.

Aralinium multiradiatum Platen.

1908 *Aralinium multiradiatum* Platen, p. 63.

Occurrence: Mio-Pliocene; California.

Aralinium parenchymaticum Platen.

1908 *Aralinium parenchymaticum* Platen, p. 62.

Occurrence: Mio-Pliocene; California.

[**Aulacoxylon** Combes, 1907, p. 28.]

[**Aulacoxylon sparnacense** Combes.]

1907 *Aulacoxylon sparnacense* Combes, p. 28, pl. I, f. 1—3.

Remarks: Neither the figures nor the very brief description give any idea of the structure of this wood, which is absolutely unidentifiable. The author does not even seem to have cut thin sections.

Occurrence: Eocene (Sparnacian); France (Auteuil).

Banksia L.

Banksia sp.

1875 *Banksia* (fossil wood): Johnston, p. 53, f. 10—13.

1888 *Banksia* (fossil wood): Johnston, pp. 270—271, pl. XLVIII, f. 8—11.

Remarks: Johnston states that the wood shows "the original structure very perfectly, and can hardly be distinguished from the sections of existing woods of the same class". The illustrations are poor and rather diagrammatic.

Occurrence: Tertiary; Tasmania (Launceston basin).

Banksioxylon Crié.

[**Banksioxylon australe** Crié.]

1889 *Banksioxylon australe* Crié, p. 78.

Remarks: Nomen nudum.

Occurrence: Pleistocene; Australia.

[Banksioxylon tasmanicum Crié.]1889 *Banksioxylon tasmanicum* Crié, pp. 90, 91.

Remarks: Nomen nudum.

Occurrence: Pliocene; Tasmania.

Betula L.(See also *Betulinium*.)**[Betula macclintockii Cramer.]**Synonym of *Betulinium macclintockii*, q. v.**[Betula salzhausensis Goepfert.]**See *Betulinium* sp.**Betula sp.**1914 *Betula* sp.: Szafer, p. 348.1926 *Betuloxylon oligocenicum* Kaiser: Fietz, p. 229.

Remarks: Fietz's record is based on a Quaternary lignite, which he claims is identical in structure with recent *Betula* wood. It is therefore difficult to understand why he uses Kaiser's name; it would seem better to describe this Silesian lignite simply as *Betula* sp., and not to confuse it with an Oligocene wood which may be specifically distinct. (See also the remarks on nomenclature in the Introduction, above.) Fietz himself is inconsistent in his terminology, for in the same paper he refers another Quaternary lignite to a living species (*Viscum album*).

Occurrence: Pleistocene; Poland, Silesia (Sörgsdorf).

Betulinium Unger, 1842, p. 101.**[Betulinium diluviale (Unger) Felix.]**Synonym of *Laurinium diluviale*, q. v.See also *Betulinium* sp.**Betulinium geinitzii (Lakowitz) Nagel.**1890 *Betuloxylon geinitzii* Lakowitz, p. 25, pl. I, f. 1-4.1916 *Betulinium geinitzii* (Lakowitz) Nagel, p. 104.

Occurrence: Oligocene; Saxony (Annaberg).

Betulinium lignitum (Kraus) Nagel.1865 *Betula lignitum* Kraus, p. 48.1890 *Betula lignitum* Kraus: Kaiser, p. 7.1916 *Betulinium lignitum* (Kraus) Nagel, p. 104.

Remarks: The wood described by Schenk (1869) and Beck (1882, 1886) as *Betula salzhausensis*, or *Betulinium* sp. (q. v.) should perhaps be included here. Kraus was of opinion that the birch woods of the Brown Coal could not be specifically distinguished.

Occurrence: Oligocene; Bavaria (Rückers, Bischofsheim, Salzhausen).

Betulinium macclintockii (Cramer) Schimper.

- 1868 *Betula macclintockii* Cramer, p. 174, pl. XXXIV, f. 4; XXXIX, f. 1—9.
 1872 *Betulinium macclintockii* (Cramer) Schimper, p. 575.
 1882 *Betulinium macclintockii* (Cramer): Felix, p. 39.
 1887 *Betula macclintockii* Cramer: Kobbe, p. 53.
 1890 *Betuloxylon macclintockii* (Cramer) Lakowitz, p. 29.
 1890 *Betula macclintockii* Cramer: Kaiser, p. 7.
 1907 *Betulinium macclintockii* (Cramer): Reiss, p. 199.
 1916 *Betulinium macclintockii* (Cramer): Nagel, p. 104.
 Occurrence: Tertiary; Banksland (Ballast Bay), Japan.

Betulinium oligocaenicum (Kaiser) Nagel.

- 1880a *Betuloxylon oligocaenicum* Kaiser, p. 511.
 1890 *Betuloxylon oligocaenicum* Kaiser: Kaiser, p. 7.
 1890 *Betuloxylon oligocaenicum* Kaiser: Lakowitz, p. 29.
 1916 *Betulinium oligocaenicum* (Kaiser) Nagel, p. 104.
 Remarks: The superfluous name *Betuloxylon* was rightly rejected by Nagel on grounds of priority. Original in Mus. Univ. Halle. Wood of this species recorded by Fietz from the Pleistocene of Silesia is here re-named *Betula* sp.
 Occurrence: Oligocene; Siebengebirge (Stein near Oberkassel).

Betulinium parisiense Unger.

- 1845 *Betulinium parisiense* Unger, p. 215.
 1847 *Betulinium parisiense* Unger: Unger, p. 119.
 1847 *Betulinium parisiense* Unger: Graves, p. 709.
 1849 *Betulinium parisiense* Unger: Unger, p. 319.
 1859 *Betulinium parisiense* Unger: Unger, p. 398.
 1857 *Betulinium parisiense* Unger: Unger, p. 11, pl. III, f. 4, 5.
 1866 *Betulinium parisiense* Unger: Watelet, p. 131.
 1872 *Betulinium parisiense* Unger: Schimper, p. 575.
 1873 *Betulinium* cf. *parisiense* Unger: Stur, p. 9.
 1882 *Betulinium parisiense* Unger: Felix, pp. 38, 40.
 1890 *Betulinium parisiense* Unger: Kaiser, p. 8.
 1890 *Betuloxylon parisiense* (Unger) Lakowitz, p. 29.
 1909 *Betuloxylon parisiense* (Unger): Fritel, p. 151.
 1916 *Betulinium parisiense* Unger: Nagel, p. 105.
 Remarks: Unger gives "*Exogenites* Brongniart" as a synonym of *Betulinium parisiense*, but this is a misuse of Brongniart's name, which was intended to have a wider significance.
 Felix (1882, p. 40) doubts whether this wood belongs to the *Betulaceae*, as also does Lakowitz (1890). Graves (1847) gives a long list of localities for this species and Fritel (1909) also gives a list of localities where fossil wood has been found in the Eocene sands of the Paris basin, but neither author states whether all of them have been examined and identified as *Betulinium*.
 Occurrence: Eocene (Cuisian); Near Paris.
 ? Tertiary; Poland (Wieliczka).

Betulinium paronae Pampaloni.

- 1904 *Betulinium paronae* Pampaloni, p. 547, f. 12—14.
 Occurrence: Pliocene; Piedmont (Astigiano).

Betulinium priscum Felix.

- 1884 *Betulinium priscum* Felix, p. 8, pl. IV, f. 2.
 1889 *Betulinium priscum* Felix: Staub, p. 186.
 1890 *Betulinium priscum* Felix: Kaiser, p. 8.
 1890 *Betuloxylon priscum* (Felix) Lakowitz, p. 29.
 1916 *Betulinium priscum* Felix: Nagel, p. 105.

Remarks: Felix names the wood *B. priscum* "nov. sp.", but states that it probably belongs to the birch leaves from the same deposit which have been named *Betula prisca* Ett. Felix discusses the differences between this wood and other species of *Betulinium*. Originals in Mus. Ungar. Geol. Reichsanst. Wood from the same locality had previously been noted by Stur, but not described.

Occurrence: Tertiary; Hungary (Medgyaszo).

Betulinium rocae (Conwentz) Nagel.

- 1885 *Betuloxylon rocae* Conwentz, p. 452.
 1890 *Betuloxylon rocae* Conwentz: Kaiser, p. 8.
 1890 *Betuloxylon rocae* Conwentz: Lakowitz, p. 29.
 1916 *Betulinium rocae* (Conwentz) Nagel, p. 105.

Remarks: A poorly preserved wood from southern Patagonia is tentatively compared by Kräusel (1924) with this species.

Occurrence: Oligocene; Argentina (Fresno-Menoco).

Betulinium rossicum Mercklin.

- 1855 *Betulinium rossicum* Mercklin, p. 33, pl. IV, f. 1, 2; V, f. 1—11.
 1865 *Betulinium rossicum* Mercklin: Eichwald, p. 57.
 1880 *Betulinium rossicum* Mercklin: Krendowsky, p. 216 [Not seen].
 1882 *Betulinium rossicum* Mercklin: Felix, p. 38.
 1890 *Betuloxylon rossicum* (Mercklin) Lakowitz, p. 29.
 1890 *Betulinium rossicum* Mercklin: Kaiser, p. 8.

Occurrence: ? Horizon; Russia (Kursk).

Betulinium stagnigenum Unger.

- 1850 *Betulinium stagnigenum* Unger, p. 398.
 1857 *Betulinium stagnigenum* Unger: Unger, p. 11, pl. III, f. 6, 7.
 1872 *Betulinium stagnigenum* Unger: Schimper, p. 575.
 1882 *Betulinium stagnigenum* Unger: Felix, p. 39.
 1890 *Betulinium stagnigenum* Unger: Kaiser, p. 9.
 1890 *Betuloxylon stagnigenum* (Unger) Lakowitz, p. 29.
 1916 *Betulinium stagnigenum* Unger: Nagel, p. 106.

Remarks: According to Lakowitz, the generic determination is doubtful.

Occurrence: Miocene (Süsswasserkalk); Bohemia (Tuchoritz).

Betulinium tenerum Unger.

- 1842 *Betulinium tenerum* Unger, p. 101.
 1842a *Betulinium tenerum* Unger: Unger, p. 173.
 1845 *Betulinium tenerum* Unger: Unger, p. 215.
 1847 *Betulinium tenerum* Unger: Unger, p. 118, pl. XXXIV, f. 8—10.
 1850 *Betulinium tenerum* Unger: Unger, p. 398.
 1852 *Betulinium tenerum* Unger: Unger, p. 106, pl. XXXIX, f. 13.
 1872 *Betulinium tenerum* Unger: Schimper, p. 575.
 1882 *Betulinium tenerum* Unger: Felix, p. 39.
 1883 *Betulinium tenerum* Unger: Hofmann, p. 88.
 1887 *Betulinium tenerum* Unger: Kobbe, p. 40.

1890 *Betulinium tenerum* Unger: Kaiser, p. 9.

1890 *Betuloxylon tenerum* (Unger) Lakowitz, p. 29.

Remarks: Originals said to be in the National Museum at Linz. Considered by Lakowitz to be of doubtful determination.

Occurrence: Tertiary (Miocene?); Austria (Freystadt, St. Stephen), Mecklenburg.

Betulinium sp.

1869 *Betula salzhausemsis* Goepfert: Schenk, p. 379.

1882 *Betulinium* sp.: Beck, p. 760.

1886 *Betula salzhausemsis*? Goepfert: Beck, p. 351.

1890 *Betula salzhausemsis* Goepfert: Kaiser, p. 8.

Remarks: Bark and poorly preserved wood, identified by Schenk (1869) with the leaf-species *Betula salzhausemsis* Goepfert, but Beck (1882) rightly considers that it is best described as *Betulinium* sp. It might perhaps be included in *Betulinium lignitum* Kraus. See also Nagel (1916, pp. 89, 90, 106—7), who wrongly ascribes the combination *Betulinium salzhausemsis* to Schenk.

Occurrence: Oligocene; Saxony.

Betulinium sp.

1882 *Betulinium diluviale* (Unger) Felix, p. 37.

1883 *Betulinium*: Felix, p. 62.

1890 *Betulinium diluviale* Felix: Kaiser, p. 7.

Remarks: Felix in 1882 united some silicified woods from Cracow with Unger's *Ulmium diluviale* under the name *Betulinium*, but later he showed that Unger's species belonged to the *Lauraceae* (see *Laurinum diluviale*). He left the Cracow wood in *Betulinium* without any specific name (1883, p. 62) but Kaiser lists it as *B. diluviale* Felix. This name cannot stand, and as the wood has never been figured, it would not be proper at present to give it a new specific name. I therefore enter it here as *Betulinium* sp.

Occurrence: Tertiary; Poland (near Cracow).

Betulinium sp.

1884 *Betulinium* sp.: Hofmann, p. 194 (43).

Remarks: Poorly preserved and not definitely identifiable.

Occurrence: Horizon and locality unknown (Utrecht Museum).

? Betulinium sp.

1873 *Betula* sp.: Dawson, p. 68.

1888 *Betula*: Dawson, p. 33.

Remarks: No figures or descriptions. Requires confirmation.

Occurrence: Cretaceous; Vancouver Island.

Upper Cretaceous (Belly River Series); Western Canada (Ribstone Creek).

Betulinium sp.

1920 *Betula* sp.: Kräusel, p. 456, pl. XXXIII, f. 6; XXXIV, f. 2—4; text-fig. 5—7.

Occurrence: Miocene; Silesia (Kohlfurt and Beuthen).

[**Betuloxylon** Kaiser, 1880, p. 511.]

Synonym of **Betulinium**, q. v. for all species described under **Betuloxylon**.

[**Bredaea** Goeppert, 1854, p. 55.]

Sunk in **Dipterocarpoxylo**, q. v. for **B. moroides**.

Brongniartites Unger, 1845, p. 264.

[Non Zalessky, 1927, Mem. Com. Geol., N. S. 176, p. 39.]

Brongniartites graecus Unger.

1845 **Brongniartites graecus** Unger, p. 264.

1850 **Brongniartites graecus** Unger: Unger, p. 524.

1852 **Brongniartites graecus** Unger: Prokesch-Osten & Unger, p. 857.

1890 **Brongniartites graecus** Unger: Kaiser, p. 34.

1898 **Brongniartites graecus** Unger: Fliche, p. 302.

Occurrence: Tertiary; Greece (Lesbos).

Bronnites Unger, 1842, p. 102.

Bronnites antiguensis Unger.

1833 **Dicotyledonous** wood: Witham, p. 67, pl. XVI, f. 11.

1842 **Bronnites antiguensis** Unger, p. 102.

1842a **Bronnites antiguensis** Unger: Unger, p. 178.

1845 **Bronnites antiguensis** Unger: Unger, p. 263.

1850 **Bronnites antiguensis** Unger: Unger, p. 524.

1890 **Bronnites antiguensis** Unger: Kaiser, p. 34.

1890 **Bronnites antiguensis** Unger: Schenk, p. 902.

Remarks: In the opinion of Schenk this wood is not identifiable.

Occurrence: Tertiary; Antigua.

Bronnites orientalis Unger.

1850 **Bronnites orientalis** Unger, p. 524.

1890 **Bronnites orientalis** Unger: Kaiser, p. 35.

Occurrence: ? Cretaceous or Tertiary; Shores of the Bosporus.

Bronnites transylvanicus Ettingshausen.

1851 **Bronnites transylvanicus** Ettingshausen in Hauer, p. 74.

1889 **Bronnites transylvanicus** Ett.: Staub, p. 190.

Occurrence: Tertiary; Transylvania.

Bronnites vindobonensis Unger.

[1845 **Bronnites viennensis** Unger, p. 263.]

1850 **Bronnites vindobonensis** Unger, p. 524.

1890 **Bronnites vindobonensis** Unger: Kaiser, p. 34.

Occurrence: Tertiary; Vienna.

Caesalpinioxylon Schenk, 1890, p. 901.

(See remarks on **Caesalpinium**).

Caesalpinioxylon nathorsti (Schuster).

1910 **Caesalpinium nathorsti** Schuster, p. 10, pl. I, f. 5, 6; II, f. 7, 8.

Remarks: Originals in Nat. Riksmus., Stockholm.

Occurrence: Tertiary; Uruguay.

Caesalpinioxylon oweni (Carruthers).

- 1870 *Nicolia oweni* Carruthers, p. 310, pl. XIV, f. 1, 2.
 1883 *Nicolia oweni* Carruthers: Schenk, pp. 19, 20.
 1884a *Nicolia minor* Hofmann, p. 485, pl. III, f. 2.
 1888 *Nicolia oweni* Carruthers: Schenk, pp. 19, 23.
 1890 *Nicolia oweni* Carruthers: Kaiser, p. 31.
 1890 *Nicolia minor* Hofmann: Kaiser, p. 35.
 1901 *Nicolia oweni* Carruthers: Blanckenhorn, pp. 102, 106.
 1910 *Caesalpinium oweni* (Carruthers) Schuster, p. 8, pl. II, f. 10, 12; III, f. 16, 17; text-fig. 1.

Remarks: Schuster showed that *Nicolia oweni* Carr. probably belonged to the *Caesalpiniae* and united with it *N. minor* Hofmann, whereas *N. aegyptiaca* Unger belonged to the *Sterculiaceae*. Schenk (1883) in discussing *N. oweni*, had already suggested a possible affinity with the *Caesalpiniae*.

Occurrence: Upper Cretaceous; Egypt (Gebel Garra near Assouan, Gebel Hefhuf in Baharia), Sudan (Dra-gara near Berber).
 Oligocene; Egypt (near Cairo).
 Middle Pliocene; Egypt (desert near Wadi Natrum).
 Tertiary; Australia (Lake Eyre), Bismarck Archipelago (Dobo).

Caesalpinioxylon palembangense Kräusel.

- 1922 *Caesalpinioxylon palembangense* Kräusel, p. 247, pl. II, f. 1; III, f. 1, 2; VII, f. 6, 11; text-fig. 21.
 1923 *Shoreoxylon palembangense* (Kräusel) den Berger, p. 144.
 1925 *Caesalpinioxylon palembangense* Kräusel: Kräusel, p. 338.
 1928 *Shoreoxylon palembangense* (Kräusel): Pfeiffer & van Heurn, p. 1007.

Remarks: Kräusel (1925, p. 340) admits the possibility that this wood may belong to the *Dipterocarpaceae*.

Occurrence: Miocene; Sumatra.

Caesalpinioxylon quirogae Schenk.

- 1890 *Caesalpinioxylon quirogae* Schenk, p. 901, f. 432.
 1929 *Caesalpinioxylon quirogae* Schenk: Chiarugi, p. 418.

Occurrence: Tertiary?; Western Sahara on the coast near Huisi Aissa.

[Caesalpinium Schleiden 1855, p. 27.]

Remarks: It is not at all clear what Schleiden intended to convey by this name, and he gives no diagnosis, figure, or description. Nevertheless Schuster (1910, p. 8) adopted Schleiden's *Caesalpinium* for woods related to the *Caesalpinaceae*, although the name *Caesalpinioxylon* had been introduced by Schenk (1890). Kräusel (1922) discarded Schleiden's name in favour of *Caesalpinioxylon*, but also overlooked Schenk's work.

[Caesalpinium tirolense Schleiden.]

- 1855 *Caesalpinium tirolense* Schleiden, p. 27.
 1890 *Caesalpinium tirolense* Schleiden: Kaiser, p. 30.

Remarks: Nomen nudum. Resinous wood, stated to resemble that of *Caesalpinia echinata*.

Occurrence: Oligocene; Tyrol (Häring).

Cantia Stopes, 1915, p. 260.**Cantia arborescens** Stopes.

1915 **Cantia arborescens** Stopes, p. 260, pls. XXVI—XXVIII; text-fig. 76—78.

Remarks: Affinities uncertain. Originals in Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Lower Cretaceous (Aptian); England (Kent).

Capparidoxylon Schenk, 1883, p. 12.**Capparidoxylon geinitzi** Schenk.

1883 **Capparidoxylon geinitzi** Schenk, p. 12, pl. I, f. 3, 4.

1890 **Capparidoxylon geinitzi** Schenk: Kaiser, p. 24.

Remarks: Compared with various species of *Capparis*.

Occurrence: ? Oligocene; near Cairo.

Carpinoxylon Vater, 1884, p. 848.**Carpinoxylon compactum** Vater.

1884 **Carpinoxylon compactum** Vater, p. 848, pl. XXIX, f. 28, 29.

1887a **Carpinoxylon compactum** Vater: Felix, p. 150.

1890 **Carpinoxylon compactum** Vater: Kaiser, p. 10.

Occurrence: Lower Senonian?; Brunswick (Helmstedt).

Carpinoxylon pfefferi Platen.

1908 **Carpinoxylon pfefferi** Platen, p. 40, pl. II, f. 1, 2.

Remarks: Schönfeld (1919, p. 8) suggests that this wood may be lauraceous.

Occurrence: Miocene; California.

Carpinoxylon vasculosum Felix.

1887 **Carpinoxylon vasculosum** Felix, p. 150, pl. XXVII, f. 4, 5.

1889 **Carpinoxylon vasculosum** Felix: Staub, p. 187.

1890 **Carpinoxylon vasculosum** Felix: Kaiser, p. 10.

Remarks: Original in Mus. Ungar. Geol. Reichsanst.

Occurrence: ? Tertiary; Hungary.

Carpinoxylon sp.

1930a **Carpinoxylon**: Schönfeld, p. 70.

Occurrence: Miocene; Saxony (Deutzen).

Carpinoxylon sp.

1920 **Carpinus** sp.: Kräusel, p. 454, pl. XXXIII, f. 5; XXXVIII, f. 6, 7; text-fig. 4.

Occurrence: Miocene; Silesia (Moys).

Carpinus L.**Carpinus** sp.

1926a **Carpinus**: Fietz, p. 417.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Cassia L.**[Cassia alata L.]**

1911a *Cassia alata* L.: Schuster, p. 246, f. 8.

Remarks: Wood referred to a living species, but it has been pointed out by Hallier (1) that *Cassia alata* is herbaceous and not a tree (2) that it is an American plant which only occurs as a weed in Java.

Occurrence: Pithecanthropus Beds; Java.

Cassioxylon Felix, 1882, p. 69.**Cassioxylon anomalum Felix.**

1882 *Cassioxylon anomalum* Felix, p. 69.

1883a *Cassioxylon anomalum* Felix: Felix, p. 15, pl. II, f. 3, 5.

1884 *Cassioxylon anomalum* Felix: Vater, p. 850.

1890 *Cassioxylon anomalum* Felix: Kaiser, p. 30.

Remarks: Resembles *Cassia speciosa*. Original in Munich.

Occurrence: ? Tertiary; probably Antigua.

Cassioxylon bartholomoei Fliche.

1888 *Cassioxylon bartholomoei* Fliche, p. 572.

Occurrence: Tertiary; Algeria (Southern Oran, between Ain Sefra and Tiout).

Cassioxylon zirkeli Felix.

1884 *Cassioxylon zirkeli* Felix, p. 32, pl. III, f. 1, 5; IV, f. 1.

1889 *Cassioxylon zirkeli* Felix: Staub, p. 190.

1890 *Cassioxylon zirkeli* Felix: Kaiser, p. 30.

Remarks: Originals in Mus. Pal. Berlin.

Occurrence: Tertiary; Hungary.

Castanopsis Spach.**Castanopsis sp.**

1925 *Castanopsis* sp.: Schönfeld, p. 19.

Occurrence: Tertiary (Brown Coal); Germany.

**Casuaroxylon Goeppert & Stache — Stache
1855, p. 42.****Casuaroxylon anglia Goeppert & Stache.**

1855 *Casuaroxylon anglia* Goeppert & Stache: Stache, p. 42.

Remarks: I have found no other references to this and the following species, which are described but not figured by Stache.

Occurrence: Locality and horizon unknown.

Casuaroxylon extorre Stache.

1855 *Casuaroxylon extorre* ? Stache, p. 42.

Occurrence: Locality and horizon unknown.

Celastrinoxylon Schenk, 1888, p. 21.

Celastrinoxylon affine Schenk.

1888 **Celastrinoxylon affine** Schenk, p. 21.

Remarks: Distinct from *Rohlfisia*, which was also referred to the *Celastraceae*.

Occurrence: Tertiary; Egypt.

[**Celtitis** Tuzson, 1909, p. 376,

Celtites Tuzson, 1911, p. 50.]

Included in **Ulminium**, q. v. for **C. kleinii**.

Cercidoxylon Platen, 1908, p. 138.

Cercidoxylon zirkeli Platen.

1908 **Cercidoxylon zirkeli** Platen, p. 139, pl. II, f. 5, 6.

Occurrence: Tertiary (? Pliocene); Nebraska (Running Water River).

[**Charpentieria** Unger, 1845, p. 262.]

Remarks: Non **Charpentieria** Gaudichaud--Beaupré, 1826.

See **Dryoxylon nivium**.

Cinnamomum L.

cf. **Cinnamomum camphora** Nees & Eberm.

1925 cf. **Cinnamomum camphora**: Schönfeld, p. 19.

Occurrence: Tertiary (Brown Coal); Germany.

Combretacinium Felix, 1894, p. 91.

Combretacinium quisqualoides Felix.

1894 **Combretacinium quisqualoides** Felix, p. 90, pl. X, f. 1a—c.

Remarks: Compared with *Quisqualis pubescens*.

Occurrence: Eocene (Sumgait series); Caucasus (Apscheron).

Constantinium Unger, 1863, p. 517.

Constantinium proteoides Unger.

1863 **Constantinium proteoides** Unger in Tehihatchef p. 517.

1866a **Constantinium proteoides** Unger: Unger, p. 322, pl. XVII, f. 1, 2.

Remarks: Referred by Unger to the *Proteaceae*.

Occurrence: Tertiary; Thrace (Lake Derkos).

Cornoxylon Conwentz, 1882, p. 157.

Cornoxylon cretaceum (Caspary).

1888 **Cornus cretacea** Caspary, p. 39.

1889 **Cornus cretacea** Caspary: Caspary, p. 23, pl. V, f. 5—8; VI, f. 1—3.

1890 *Cornus cretacea* Caspary: Kaiser, p. 28.

Remarks: Compared with *Cornus alba*. Original in Mus. Bot. Gart., Königsberg.

Occurrence: Probably Cretaceous; Königsberg?.

Cornoxyton erraticum Conwentz.

1882 *Cornoxyton erraticum* Conwentz, p. 157.

1889 *Cornoxyton erraticum* Conwentz: Caspary, p. 33.

1890 *Cornoxyton erraticum* Conwentz: Kaiser, p. 29.

Remarks: A Senonian wood compared with this species by Vater was considered by Caspary to be distinct (See *Cornoxyton vateri*).

Occurrence: Pleistocene (erratic derived from an earlier formation); Holstein.

Cornoxyton holsatiae Conwentz.

1882 *Cornoxyton holsatiae* Conwentz, p. 160.

1883 *Cornoxyton holsatiae* Conwentz: Hofmann, p. 89.

1889 *Cornoxyton* "holsaticum" Conwentz: Caspary, p. 33.

1890 *Cornoxyton holsatiae* Conwentz: Kaiser, p. 29.

Occurrence: Pleistocene (derived); Mecklenburg and Holstein.

Cornoxyton latiporosum Kräusel & Schönfeld.

1924 *Cornoxyton latiporosum* Kräusel & Schönfeld, p. 277, text-fig. 20—26.

1924 *Cornoxyton* sp. (*latiporosum*?): Kräusel & Schönfeld, p. 280.

Remarks: The second entry refers to wood which, though poorly preserved, the authors consider to be probably identical with *C. latiporosum*. The species is compared with *Cornoxyton holsatiae* Conwentz.

Occurrence: Miocene; Holland (S. Limburg).

Cornoxyton myricaeforme Vater.

1884 *Cornoxyton myricaeforme* Vater, p. 846, pl. XXIX, f. 25, 26.

1890 *Cornoxyton myricaeforme* Vater: Kaiser, p. 29.

Remarks: A resemblance to *Myrica* is noted by Vater.

Occurrence: Lower Senonian; Brunswick (Helmstedt).

Cornoxyton solidior (Caspary).

1888 *Cornus cretacea* forma *solidior* Caspary, p. 40.

1889 *Cornus cretacea* forma *solidior* Caspary: Caspary, p. 29, pl. VI, f. 4—17.

1890 *Cornus cretacea* forma *solidior* Caspary: Kaiser, p. 29.

Remarks: Failing a re-investigation of the material, (the originals are in the Mus. Bot. Gart., Königsberg), the name *Cornoxyton solidior* should be used, as it is absurd to introduce a "variety" or "forma" for a fossil wood.

Occurrence: ? Tertiary; Prussia (Herzogsacker near Königsberg).

Cornoxyton vateri (Caspary).

1884 *Cornoxyton* cf. *erraticum* Conwentz: Vater, p. 846, pl. XXIX, f. 27.

1888 *Cornus vateri* Caspary, p. 40.

Remarks: See also Caspary, 1889, p. 33.

Occurrence: Lower Senonian; Brunswick (Helmstedt).

Cornoxyylon sp.

1924 *Cornoxyylon* sp.: Kräusel & Schönfeld, p. 232.

Remarks: Referred provisionally to *Cornoxyylon*. May be related to *Hamamelis* and *Liquidambar*.

Occurrence: Miocene; Holland (S. Limburg).

Corylus L.**Corylus sp.**

1926a *Corylus*: Fietz, p. 417.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Cottaites Unger, 1842, p. 101.

[*Cottaites lapidariorum* Unger.]

Synonym of *Ulminium lapidariorum* q. v.

Cottaites robustior Unger.

1842 *Cottaites robustior* Unger, p. 102.

1842a *Cottaites robustior* Unger: Unger, p. 176.

1845 *Cottaites robustior* Unger: Unger, p. 265.

1850 *Cottaites robustior* Unger: Unger, p. 526.

1889 *Cottaites robustior* Unger: Staub, p. 191.

1890 *Cottaites robustior* Unger: Kaiser, p. 32.

Remarks: The genus *Cottaites* was placed in the *Leguminosae* by Unger. *C. lapidariorum* however has since been transferred to *Ulminium*, and Schenk remarks (1890, p. 902) that the other two species are certainly not leguminous.

Occurrence: Tertiary; Hungary (Antal near Schemnitz).

Cottaites vasculosus Unger.

1845 *Cottaites vasculosus* Unger, p. 265.

1850 *Cottaites vasculosus* Unger: Unger, p. 526.

1890 *Cottaites vasculosus* Unger: Kaiser, p. 32.

Remarks: See *C. robustior*.

Occurrence: Tertiary; Austria (Nicolaiaburg).

Dicotyledonous Wood.

1835 *Dicotyledonous* wood: Nicol, pp. 335—336.

Remarks: Nicol describes, but does not name or figure, woods from Mull, Scotland, and from Egypt and Nubia. Nicol's collection is now in the Geological Department of the British Museum (Nat. Hist.); two sections of dicotyledons from Mull belong to *Plataninum*, and among a few sections from Egypt, mostly poorly preserved, one at least is *Nicolia aegyptiaca*.

Dicotyledonous Wood.

1837 *Dicotyledonous* wood: Stokes, p. 208, pl. XVII, f. 1, 2.

Occurrence: Tertiary; Antigua.

Dicotyledonous Wood.

- 1850 **Dicotyledonous** wood: Dixon, p. 235, pl. XVI, f. 6.
 1878 **Dicotyledonous** wood: Carruthers, p. 164, pl. XVI, f. 6.
 Occurrence: Eocene; England (Sussex).

Dicotyledonous Wood.

- 1852 **Dicotyledonous** wood: Sorby, p. 91, pl. XVI.
 Remarks: Said to be from Lias at Keynsham, but it was only obtained from a dealer and there is probably a mistake in the locality and horizon.

Dicotyledonous Wood.

- 1856 **Dicotyledonous** wood: Bailey, p. 337, pl. XII, f. 1, 2.
 Occurrence: ? Tertiary; U. S. A. (W. slope of Sierra Nevada).

Dicotyledonous Wood.

- 1856 **Dicotyledonous** wood: Schaeffer, p. 338, pl. XII, f. 3, 4.
 Occurrence: ? Tertiary; U. S. A. (Colorado Desert).

Dicotyledonous Wood.

- 1875 **Dicotyledonous** wood: Johnston, p. 53, f. 14, 15.
 1888 **Dicotyledonous** wood: Johnston, pp. 270, 271, pl. XLVII, f. 9, 10;
 ? pl. XLVIII, f. 6, 7.
 Remarks: The figures are poor and quite unidentifiable. Johnston mentions "two or three undetermined exogens" in addition to *Banksia* (q. v.).
 Occurrence: Tertiary; Tasmania.

Dicotyledonous Wood.

- 1882a **Dicotyledonous** wood (Cedrelaceae): Kraus, p. 6.
 Remarks: Regarded as certainly belonging to the *Cedrelaceae* (= *Meliaceae*), but not generically identifiable. Popularly supposed to be wood of a living plant which becomes silicified when buried in the soil.
 Occurrence: ? Tertiary; Mexico.

Dicotyledonous Wood.

- 1885 „Eigentümliches Laubholz": Quenstedt, p. 1151, pl. XCVIII, f. 24.
 Remarks: Quite unrecognizable from the figures given.
 Occurrence: Tertiary; Germany (Dietenheim).

Dicotyledonous Wood.

- 1902 **Dicotyledonous** wood: Knowlton, p. 640, pl. XXV, f. 14.
 1906 **Dicotyledonous** wood: Jeffrey & Chrysler, p. 199, pl. L, f. 7-12;
 LI, f. 13-15.
 Remarks: Knowlton compared this wood with *Betula*. Jeffrey & Chrysler name it *Betuloxylon* in the legend to the plate, but in the text they say "it can scarcely be a *Betuloxylon*".
 Occurrence: Eocene; Vermont (Brandon).

Dicotyledonous Wood.

1911 **Dicotyledonous** wood: Tuzson, p. 53, pl. II, f. 5; text-fig. 26—30.

1911 **Dicotyledonous** wood: Tuzson, p. 55, text-fig. 31—37.

Occurrence: Tertiary; Hungary (Kislöd and Pét).

Dicotyledonous Wood.

1912 **Dicotyledonous** wood: Compter, pp. 413—420.

Remarks: Compter describes ten different petrified specimens from a diluvial deposit, but does not name any of them. Some are compared with *Quercinium*, and one with *Populus*. Most of them seem to have been poorly preserved.

Occurrence: ?Tertiary; Weimar (Apolda).

Dicotyledonous Wood.

1914 **Dicotyledonous** wood: Oswald, p. 130.

Remarks: Calcified trunks belonging to at least four different types, perhaps referable (according to Dr. H. Bancroft) to the *Leguminosae*, the *Lauraceae* or the *Bombacaceae*, the *Humiriaceae* and the *Caprifoliaceae*. A new description of these and other African woods will shortly be published by Dr. Bancroft in the *Annals of Botany*.

Occurrence: Miocene; Kenya Colony (Kikongo near Karungu).

Dicotyledonous Wood.

1920 **Dicotyledonous** wood: Sahni, p. 34, pl. III, f. 16.

Occurrence: Tertiary; Queensland (Condamine River).

Dicotyledonous Wood.

1924 **Dicotyledonous** wood: Kräusel, p. 2, pl. I, f. 1—7.

Remarks: Perhaps belongs to the *Ericaceae* or the *Myrtaceae*.

Occurrence: Pleistocene; Chiloe.

Dicotyledonous Wood.

1924 **Dicotyledonous** wood: Kräusel, p. 28.

Remarks: Poorly preserved; perhaps close to *Betulinium rocae* (Conwentz).

Occurrence: Tertiary; Patagonia.

Dicotyledonous Wood.

1924 ? **Betulaceae**: Seward & Holtum, p. 81.

Occurrence: Lower Tertiary (?Eocene); Scotland (Mull).

Dicotyledonous Wood.

1926 [Fossilholz]: Udluft, p. 1, pl. II, f. 6—11, 14.

Occurrence: Volcanic tuffs of Mt. Elgon; Central Africa.

Dicotyledonous Wood.

1927 **Dicotyledonous** wood: Barnard, p. 113, pls. V, VI.

Remarks: The affinities are considered to be with the *Saxifragaceae*.

Occurrence: Tertiary; New South Wales (Ulladulla).

Fossilium Catalogus II. 17.

Dicotyledonous Wood.

- 1928 **Dicotyledonous** wood: Chhibber, p. 13, pl. I.
Occurrence: Late Tertiary; Burma.

Dicotyledonous Wood.

- 1928 **Dicotyledonous** wood: Donath, p. 57, f. 1—3.
Remarks: Silicified wood, perhaps oak, embedded in basalt.
Occurrence: Tertiary?; Saxony (Ostritz).

Diospyros L.**Diospyros sp.**

- 1925 **Diospyros** sp.: Schönfeld, p. 19.
Remarks: Compare Beck's record of *Ebenoxyylon tenax*.
Occurrence: Tertiary (Brown Coal); Germany.

Dipterocarpoxyylon Holden, 1916, p. 271.

Remarks: Instituted for fossil woods of the family *Dipterocarpaceae*, but used by den Berger (1927) in a more restricted sense for woods resembling those of *Dipterocarpus* and *Anisoptera* only. See also Pfeiffer and van Heurn (1928). — The wider usage has been adopted in this Catalogue.

Dipterocarpoxyylon annamense Colani.

- 1919 **Dipterocarpoxyylon annamense** Colani, p. 2, pls. I, II.
Remarks: The wood does not seem to be very well preserved, and the generic identification should be accepted with reserve.
Occurrence: ? Tertiary; Annam.

Dipterocarpoxyylon burmense Holden.

- 1916 **Dipterocarpoxyylon burmense** Holden, p. 271, pl. XXIX, f. 1—5.
1922 **Dipterocarpoxyylon burmense** Holden: Kräusel, p. 266.
1922a **Dipterocarpoxyylon burmense** Holden: Kräusel, p. 14.
1928 **Dipterocarpoxyylon burmense** Holden: Chhibber, p. 22.
Remarks: Kräusel considers that the wood is related to *Hopea* or *Shorea* rather than to *Dipterocarpus*.
Occurrence: Tertiary; Burma.

Dipterocarpoxyylon djambiense (den Berger).

- 1922 **Dipterocarpoxyylon** sp.: Kräusel, p. 269, pl. I, f. 6; II, f. 7—9; V, f. 9; VI, f. 1; VII, f. 2; text-fig. 26.
1922a **Dipterocarpoxyylon** sp.: Kräusel, p. 14.
1923 **Shoreoxyylon djambiense** den Berger, p. 147.
Remarks: See remarks on the genus *Shoreoxyylon*. On the age of the Sumatra and Java woods see van Heurn (1927).
Occurrence: Tertiary; South Sumatra.

Dipterocarpoxyylon goepperti Kräusel.

- 1926 **Dipterocarpoxyylon goepperti** Kräusel, p. 4, pl. I, f. 3, 4; II, f. 5, 6.
1927 **Dipterocarpoxyylon goepperti** Kräusel: den Berger, p. 498.
Occurrence: Tertiary; Java (Nangoeng).

Dipteroxylon javanense Kräusel.

- 1922a *Dipteroxylon javanense* Kräusel, p. 13, pl. I, f. 1—5; text-fig. 1, 2.
 1926 *Dipteroxylon javanense* Kräusel: Kräusel, p. 3.
 1927 *Dryobalanoxylon javanense* (Kräusel) den Berger, p. 498.
 Occurrence: Tertiary; Java (Bolang).

Dipteroxylon kräuseli (den Berger).

- 1922 *Dipteroxylon* sp. (*tobleri* ?): Kräusel, p. 267, pl. I, f. 4; III, f. 4, 5; VII, f. 1; text-fig. 25.
 1922a *Dipteroxylon* sp. (*tobleri* ?): Kräusel, p. 14.
 1923 *Shoreoxylon kräuseli* den Berger, p. 147.
 Remarks: See remarks on the genus *Shoreoxylon*.
 Occurrence: Tertiary; South Sumatra.

Dipteroxylon moroides (Goeppert) Kräusel.

- 1854 *Bredaea moroides* Goeppert, p. 56, pl. I, f. 3—5.
 1890 *Bredaea moroides* Goeppert: Kaiser, p. 18.
 1926 *Dipteroxylon moroides* (Goeppert) Kräusel, p. 4, pl. I, f. 2; II, f. 3, 4.
 1927 *Shoreoxylon moroides* (Goeppert) den Berger, p. 498.
 Occurrence: Tertiary; Java.

Dipteroxylon porosum (Stopes) Kräusel.

- 1912 *Woburnia porosa* Stopes, p. 9, pl. VII, f. 7; VIII, f. 8; text-fig. 6.
 1915 *Woburnia porosa* Stopes: Stopes, p. 267, text-fig. 79—81.
 1922 *Woburnia "scottii"* Stopes: Kräusel, p. 266.
 1922a *Dipteroxylon "scottii"* Stopes: Kräusel, pp. 12, 14.
 Remarks: See remarks on the name *Woburnia*. Stopes stated that the wood was "in complete agreement with some of the *Dipterocarpaceae*", and Kräusel transferred it to the form-genus *Dipteroxylon*, though he made an obvious mistake in quoting the specific name. Originals in Geol. Dept., Brit. Mus. (Nat. Hist.). Also figured Scott, D. H. 1924. *Extinct Plants*, p. 54, f. 6.
 Occurrence: Lower Cretaceous (Aptian); England (Woburn).

Dipteroxylon spectabile (Crié) Kräusel.

- 1888 *Naucleoxylon spectabile* Crié, p. 19, pl. VIII, f. 1, 2.
 1926 *Dipteroxylon spectabile* (Crié) Kräusel, p. 2, pl. I, f. 1; II, f. 1, 2.
 1927 *Dryobalanoxylon spectabile* (Crié) den Berger, p. 498.
 Occurrence: Tertiary; Java.

Dipteroxylon swedenborgii (Schuster) Kräusel.

- 1910 *Grewioxylon swedenborgii* Schuster, p. 14, pl. I, f. 1—4; text-fig. 3.
 1922 *Dipteroxylon swedenborgii* (Schuster): Kräusel, p. 267.
 1922a *Dipteroxylon swedenborgii* (Schuster): Kräusel, pp. 12, 14.
 Remarks: Kräusel showed that this wood had nothing to do with *Grewia*, but was close to his *Dipteroxylon tobleri*.
 Occurrence: Tertiary; East Indies.

Dipterocarpoxylo. tobleri Kräusel.

- 1922 **Dipterocarpoxylo. tobleri** Kräusel, p. 263, pl. I, f. 5; II, f. 6; III, f. 3; VI, f. 8; VII, f. 10; text-fig. 24.
1922a **Dipterocarpoxylo. tobleri** Kräusel: Kräusel, p. 14.
1923 **Dryobalanoxylo. tobleri** (Kräusel) den Berger, p. 146.
1925 **Dipterocarpoxylo. tobleri** Kräusel: Kräusel, p. 340.
1928 **Dryobalanoxylo. tobleri** (Kräusel): Pfeiffer & van Heurn, p. 1007.
Occurrence: Tertiary; South Sumatra.

Dipterocarpoxylo. spp.

- 1927 **Dryobalanoxylo. sp.**: van Heurn, p. 284, pl. I, f. 2, 3.
1927 **Shoreoxylo. sp.?**: van Heurn, p. 284, pl. I, f. 4.
1928 **Dipterocarpoxylo., Dryobalanoxylo., Shoreoxylo. spp.**: Pfeiffer & van Heurn, p. 1006.
Remarks: Pfeiffer & van Heurn examined a large series of woods from Bolang, three of which were figured by van Heurn (1927), but they say that "no attempt was made to establish which of the species found of this family [*Dipterocarpaceae*] corresponded with the species previously described by Kräusel and den Berger."
According to van Heurn (1927, p. 285) the fossil woods of Sumatra belong in the main to the Pliocene, and those of Java to the Plio-Pleistocene.
Occurrence: Plio-Pleistocene; Java (Bolang).

? Dipterocarpoxylo. sp.

- 1922 **? Dipterocarpoxylo. sp.**: Kräusel, p. 271, pl. I, f. 7; IV, f. 1; text-fig. 27.
Remarks: For other woods first described by Kräusel as *D. sp.*, see *D. djambiense* and *D. kräuseli*.
Occurrence: Tertiary; South Sumatra.

Djambioxylo. Kräusel, 1922, p. 272.

Djambioxylo. sumatrense Kräusel.

- 1922 **Djambioxylo. sumatrense** Kräusel, p. 272, pl. II, f. 2; IV, f. 7; V, f. 1, 6—8; VII, f. 7—9, 12; text-fig. 28.
1923 **Djambioxylo. sumatrense** Kräusel: den Berger, p. 147.
Remarks: Perhaps belongs to the *Sapindaceae*.
Occurrence: Tertiary; Sumatra.

Dombeyoxylo. Schenk, 1883, p. 13.

Dombeyoxylo. aegyptiacum Schenk.

- 1883 **Dombeyoxylo. aegyptiacum** Schenk, p. 13.
1887 **Dombeyoxylo. aegyptiacum** Schenk: Felix, p. 522.
1890 **Dombeyoxylo. aegyptiacum** Schenk: Kaiser, p. 24.
1910 **Dombeyoxylo. aegyptiacum** Schenk: Schuster, p. 12, pl. III, f. 18.
Remarks: Compared by Schenk with the wood of the *Sterculiaceae*, and especially with the recent genera *Ruizia* and *Guazuma*; by Felix with *Guazuma*; and by Schuster with *Eriodendron*. Original in Mus. Pal. Dresden.
Occurrence: ? Oligocene; Egypt (Seeber near Tureh; between Gart el Leben and Moghara).

Dombeyoxylon affine Felix.1887 **Dombeyoxylon affine** Felix, p. 520, pl. XXV, f. 2, 3, 5.1890 **Dombeyoxylon affine** Felix: Kaiser, p. 24.Remarks: Compared with the recent *Dombeya mollis* (*Sterculiaceae*). Originals in Min. Mus. Berlin.

Occurrence: Tertiary; Abyssinia (Edda Jesus near Axum).

Dombeyoxylon jacksonensis Berry.1924 **Dombeyoxylon jacksonensis** Berry, p. 181, pls. XXXVI, XXXVII.Remarks: Compared with *D. affine* Felix from Abyssinia. Type in U. S. Nat. Mus.

Occurrence: Eocene (Fayette sandstone); Louisiana (Hornbeck).

[Dryobalanoxylon den Berger, 1923, p. 146 & 1927, p. 497.]For dipterocarpaceous woods resembling *Dryobalanops*. Here included (following Kräusel) in *Dipterocarpoxyylon*.**Dryoxylon** Schleiden in Schmid, 1853, p. 28.

Remarks: This form-genus was expressly founded for dicotyledonous woods of uncertain affinity, but unfortunately it seems to have been entirely overlooked. The name *Exogenites* was also founded for a similar purpose by Brongniart in 1822, but as in those days the term "dicotyledon" included conifers and even tree lycopods, Schleiden's name is preferable, for he says "Ich zog daher den allgemeinen Namen *Dryoxylon* Holz eines Laubholz-Baumes vor". As it happens, the only species described by Schleiden was probably not a dicotyledon at all, but that does not affect the use of a form-genus name.

A score or more of Unger's species, to mention no others, might well be listed merely as *Dryoxylon*, but it has not seemed worth while to make the transfer, when one has not had an opportunity of examining the original specimens, except in the cases where Unger's generic names were pre-occupied. I have also included in *Dryoxylon* a few species of somewhat doubtfully identified woods (such as *Erica sambiensis*) in order to avoid the unwarrantable use of a recent generic name on the one hand, or the institution of yet another form-name on the other.

Dryoxylon americanum (Unger).1852a **Roemeria americana** Unger, p. 95.Remarks: The name *Roemeria* is pre-occupied.

Occurrence: Cretaceous; Texas.

[Dryoxylon jenense Schleiden.]1853 **Dryoxylon jenense** Schleiden in Schmid, p. 28.Remarks: A fragmentary wood compared with that of *Salix*, which can hardly be correct if the fossil really came from the Trias. Either the identification of the wood or the horizon must be wrong.

Occurrence: Lower Muschelkalk; Germany (Wogau near Jena).

Dryoxylon laxum (Caspary).1888 *Magnolia laxa* Caspary, p. 38.1889 *Magnolia laxa* Caspary: Caspary, p. 1, pl. I, f. 1—12.1890 *Magnolia laxa* Caspary: Kaiser, p. 24.Remarks: The identification with *Magnolia* does not seem absolutely certain, and I have therefore transferred the species to *Dryoxylon*.

Occurrence: Tertiary; East Prussia.

Dryoxylon nivium (Unger).1845 *Charpentiera nivium* Unger, p. 262.1850 *Charpentiera nivium* Unger: Unger, p. 523.1890 *Charpentiera nivium* Unger: Kaiser, p. 35.Remarks: The name *Charpentiera* is pre-occupied.

Occurrence: ?Tertiary; Galicia.

Dryoxylon sambiense (Caspary).1888 *Erica sambiensis* Caspary, p. 41.1889 *Erica sambiensis* Caspary: Caspary, p. 34, pl. VI, f. 18; VII, f. 1—10.1890 *Erica sambiensis* Caspary: Kaiser, p. 33.Remarks: Original in Mus. Bot. Gart., Königsberg. Very similar to *E. vagans*, but also compared with some *Myrtaceae*.

Occurrence: Oligocene; Prussia (Palmnicken).

Dryoxylon silvaticum (Tuzson).1909 *Magnolites silvatica* Tuzson, p. 376 (Nomen).1911 *Magnolites silvatica* Tuzson: Tuzson, p. 44, pl. II, f. 4; text-fig. 17—21.Remarks: The form-genus *Magnolites* should not be used for fossil woods.

Occurrence: Tertiary (Schotter Beds); Hungary (Lake Balaton).

Ebenoxylon Felix, 1882, p. 71.**Ebenoxylon boreale** Platen.1908 *Ebenoxylon boreale* Platen, p. 147.

Occurrence: Tertiary (? Oligocene); Alaska.

Ebenoxylon diospyroides Felix.1882 *Ebenoxylon diospyroides* Felix, p. 71, pl. I, f. 3.1883a *Ebenoxylon diospyroides* Felix: Felix, p. 17, pl. IV, f. 6.1890 *Ebenoxylon diospyroides* Felix: Kaiser, p. 33.Remarks: Similar to wood of *Diospyros*, especially to *D. discolor* and *D. virginiana*.

Occurrence: Tertiary; Antigua.

Ebenoxylon ebenoides (Schenk).1880 *Jordania ebenoides* Schenk, p. 660.1883 *Jordania ebenoides* Schenk: Schenk, p. 10, pl. IV, f. 13, 14.1890 *Jordania ebenoides* Schenk: Kaiser, p. 34.Remarks: The generic name *Jordania* is pre-occupied. The wood is compared with that of the *Ebenaceae*.

Occurrence: Upper Cretaceous; Libyan desert (near Regenföld).

Ebenoxylon speciosum Platen.

- 1908 *Ebenoxylon speciosum* Platen, p. 68.
Occurrence: Tertiary; California.

Ebenoxylon tenax Beck.

- 1886 *Ebenoxylon tenax* Beck, p. 348, pl. VII, f. 7—9.
1890 *Ebenoxylon tenax* Beck: Kaiser, p. 34.
1930a *Ebenoxylon tenax* Beck: Schönfeld, p. 70.
Remarks: Compared with wood of *Diospyros ebenus*. Schönfeld (1925) refers to Beck's record as wood of *Diospyros*.
Occurrence: Oligocene; Saxony.

Ebenoxylon tunetana (Fliche).

- 1888 *Jordania tunetana* Fliche, p. 571.
1893 *Jordania tunetana* Fliche: Thomas, p. 3.
Remarks: Compared with *E. ebenoides* (Schenk).
Occurrence: Pliocene; Tunisia (Ain Cherichira).

Ebenoxylon sp.

- 1898 *Ebenoxylon* sp.: Fliche, p. 298.
Occurrence: ? Tertiary; Mytilene (Orthymnos).

Elaeodendroxylon Platen, 1908, p. 120.

Elaeodendroxylon polymorphum Platen.

- 1908 *Elaeodendroxylon polymorphum* Platen, p. 120.
Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

[**Erica sambiensis** Caspary.]

See *Dryoxylon sambiense*.

Eucalyptus L'Hérit.

Eucalyptus sp.

- 1918 *Eucalyptus* sp. aff. *melliodora* Cunningham: Chapman, p. 172, pl. X.
1918 *Eucalyptus* sp. aff. *piperita* Smith: Chapman, p. 174.
Occurrence: Tertiary; Victoria (Gippsland).

Eugenia Mich.

Eugenia cordata Laws.

- 1912 *Eugenia cordata* Laws.: Warren, p. 367, pls. XXVI, XXVII.
Remarks: Lignitic and partly silicified wood of late date, identified with a Myrtaceous tree which still grows in the vicinity.
Occurrence: Late Tertiary or Pleistocene; coast of Zululand.

Euonyminium Mercklin, 1855, p. 23.**Euonyminium auerbachii** Mercklin.1855 **Euonyminium auerbachii** Mercklin, p. 23, pl. I, f. 3; III, f. 1—11.1865 **Euonyminium auerbachii** Mercklin: Eichwald, p. 66.1890 **Euonyminium auerbachii** Mercklin: Kaiser, p. 27.Remarks: Compared with *Euonymus japonicus* by Mercklin. Eichwald doubts the attribution, and even suggests that it may be a conifer.

Occurrence: ? Cretaceous; Russia (Durasovka in Saratov).

Euonymus L.**Euonymus** sp.1926a **Euonymus**: Fietz, p. 418.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Euphorbioxylon Felix, 1887, p. 525.**Euphorbioxylon speciosum** Felix.1887 **Euphorbioxylon speciosum** Felix, p. 525, pl. XXV, f. 4, 6, 7.1890 **Euphorbioxylon speciosum** Felix: Kaiser, p. 28.Remarks: Compared by Felix with various genera of the *Euphorbiaceae*, including *Euphorbia*, *Jatropha*, and *Hippomane*.

Occurrence: ? Tertiary; Columbia (Sabanilla).

[Exogenites Brongniart, 1822, p. 354.]

Founded as a form-genus for fossil dicotyledonous wood (but see remarks on *Dryoxylon*). No species mentioned. Unger includes *Exogenites* as a synonym of *Betulinium parisiense*, which is obviously unwarrantable. Graves (1847, p. 710) gives a list of localities where "*Exogenites*" had been found in the department of Oise, but there are no descriptions, and some of the wood was probably coniferous.

[Fagoxylon Stopes & Fujii 1910.]Synonym of **Fegonium**.**Fagus** L.**Fagus** sp.1925 **Fagus** sp.: Schönfeld, p. 19.

Occurrence: Tertiary (Brown Coal); Germany.

Fegonium Unger (1842, p. 101) em. Vater.

Vater (1884, p. 836) gives a diagnosis of *Fegonium* "gen. nov." and later explains that the diagnosis given by Unger (1847, p. 103; 1850, p. 407) for *Fegonium* is almost identical with that given for *Plataninium* (1847, p. 138; 1850, p. 414) and further that the species of *Fegonium* described by Unger belong in fact to *Plataninium*.

The spelling *Phegonium* appears in Unger's earliest works.*Fagoxylon* Stopes & Fujii 1910 should be included in *Fegonium*.

Fegonium caucasicum Felix.1894 *Fegonium caucasicum* Felix, p. 102, pl. VIII, f. 2.Remarks: Felix compares and contrasts this wood with other species of *Fegonium*.

Occurrence: Eocene (Sumgait Series); Caucasus (Apscheron).

Fegonium dryandraeforme Vater.1884 *Fegonium dryandraeforme* Vater, p. 838, pl. XXVIII, f. 7—10.1890 *Fegonium dryandraeforme* Vater: Kaiser, p. 10.

Occurrence: Lower Senonian (derived from); Brunswick.

Fegonium hokkaidense (Stopes & Fujii).1910 *Fagoxylon hokkaidense* Stopes & Fujii, p. 64, pl. VIII, f. 50—53.

Remarks: Originals in Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Upper Cretaceous; Japan (Hokkaido).

Fegonium lignitum Beck.1886 *Fegonium lignitum* Beck, p. 350.1890 *Fegonium lignitum* Beck: Kaiser, p. 10.1930a *Fegonium lignitum* Beck: Schönfeld, p. 70.Remarks: Said to resemble *F. dryandraeforme* Vater.

Occurrence: ? Oligocene; Saxony (Raupenhaim, Deutzen).

[Fegonium megapolitanum Hofmann.]Synonym of *Plataninium megapolitanum*, q. v.**[Fegonium salinarum** Unger.]Synonym of *Plataninium salinarum*, q. v.**Fegonium schenki** Vater.1884 *Fegonium schenki* Vater, p. 839, pl. XXVIII, f. 11—14.1890 *Fegonium schenki* Vater: Kaiser, p. 11.

Occurrence: Lower Senonian (derived from); Brunswick.

[Fegonium vasculosum Unger.]Synonym of *Plataninium vasculosum*, q. v.**Felixia** Platen, 1908, p. 66.**Felixia latiradiata** Platen.1908 *Felixia latiradiata* Platen, p. 66, pl. II, f. 3, 4.Remarks: The name *Felixia* was proposed for leguminous woods which cannot be more nearly identified.

Occurrence: Mio-Pliocene; California.

Fichtelites Unger, 1842, p. 101.**Fichtelites articulatus** Unger.

- 1842 **Fichtelites articulatus** Unger, p. 101.
 1842a **Fichtelites articulatus** Unger: Unger, p. 175.
 1842b „**Fichtelia articulata**” Unger: Unger, p. 748.
 1845 **Fichtelites articulatus** Unger: Unger, p. 264.
 1850 **Fichtelites articulatus** Unger: Unger, p. 525.
 1890 **Fichtelites articulatus** Unger: Kaiser, p. 32.

Remarks: Put in *Leguminosae* by Unger. Originals said to be in Nat. Mus. Linz.

Occurrence: Tertiary; Upper Austria.

Ficoxylon Kaiser, 1880, p. 309.[**Ficoxylon bohemicum** Kaiser.]

Synonym of **F. tropicum**, q. v.

Ficoxylon cretaceum Schenk.

- 1883 **Ficoxylon cretaceum** Schenk, p. 14, pl. V, f. 17—19.
 1888 **Ficoxylon cretaceum** Schenk: Fliche, p. 571.
 1890 **Ficoxylon cretaceum** Schenk: Kaiser, p. 17.
 1893 **Ficoxylon cretaceum** Schenk: Thomas, p. 3.
 1901 **Ficoxylon schenki** Blanckenhorn, p. 113.
 1929a **Ficoxylon cretaceum** Schenk: Chiarugi, p. 561.

Remarks: *F. cretaceum* Schenk was re-named by Blanckenhorn because it was definitely not of Cretaceous age. This is not a sufficient reason for disregarding the rule of priority; moreover, wood of this type may yet be found in the upper Cretaceous. Compared by Schenk with *Ficus sycomorus*.

Occurrence: Oligocene; Egypt.

Tertiary; Tunisia (Ain Cherichira), Sirtica.

Ficoxylon helictoxyloides Platen.

- 1908 **Ficoxylon helictoxyloides** Platen, p. 51.
 Occurrence: Mio-Pliocene; California.

[**Ficoxylon schenki** Blanckenhorn.]

Synonym of **F. cretaceum**, q. v.

Ficoxylon tropicum (Schleiden).

- 1855 **Ungerites tropicus** Schleiden, p. 37.
 1880 **Ficoxylon bohemicum** Kaiser, p. 309.
 1883 **Ficoxylon tropicum** (Schleiden) Felix, p. 81.
 1890 **Ficoxylon tropicum** (Schleiden): Kaiser, p. 17.

Remarks: According to Felix, the wood is close to that of *Ficus cordata*.

Occurrence: Tertiary; Bohemia (Kostenblatt and Zettow).

Ficoxylon zirkeli Hofmann.1884 **Ficoxylon zirkeli** Hofmann, p. 185 (34).1890 **Ficoxylon zirkeli** Hofmann: Kaiser, p. 17.Remarks: Silicified wood, comparable with *Ficus elastica*. Original in Coll. Univ. Utrecht.

Occurrence: Locality and horizon unknown (? Coburg).

Ficus L.**Ficus callosa** Willd.1911a **Ficus callosa** Willd.: Schuster, p. 246, f. 6.

Remarks: Wood referred to a living species. The determination should be accepted with reserve.

Occurrence: Pithecanthropus Beds; Java.

Fraasia Unger, 1850, p. 457.**Fraasia sapindoides** Unger.1850 **Fraasia sapindoides** Unger, p. 457.1852 **Fraasia sapindoides** Unger: Massalongo, p. 24.1890 **Fraasia sapindoides** Unger: Kaiser, p. 26.Remarks: Placed in the *Sapindaceae* by Unger.

Occurrence: Tertiary; probably Hungary.

[Fraxinoxylon Hofmann, 1929, p. 82.]Remarks: For Oleaceous woods resembling *Fraxinus*, the name *Ornoxylon* Felix has priority. Hofmann speaks of *Fraxinoxylon excelsior*, which is an impossible combination.**Fraxinus** L.(See also **Ornoxylon**.)**Fraxinus** cf. **excelsior** L.1882 **Fraxinus** cf. **excelsior** L.: Conwentz, p. 147.1923 **Fraxinus excelsior** Hofmann, p. 1, pl. 1, f. 1.1929 **Fraxinoxylon excelsior** Hofmann, p. 82.Remarks: If the wood mentioned by Hofmann is identical with that of *Fraxinus excelsior*, there is no point in using the name *Fraxinoxylon*, which in any case would give way to the pre-existing *Ornoxylon*. And if a form-name is to be used, the specific name *excelsior* is inadmissible. The wood may be called *Fraxinus* cf. *excelsior*.

Occurrence: Pleistocene; Holstein.

Plio-Pleistocene; Hungary (Csadberge).

Fraxinus sp.1914 **Fraxinus** sp.: Szafer, p. 349.

Occurrence: Pleistocene; Poland.

Grewioxylon Schuster, 1910, p. 14.

Remarks: The only species has been transferred by Kräusel to *Dipterocarporoxylon* (See *D. swedenborgii*).

Hamamelidoxylon Lignier, 1907, p. 300.**Hamamelidoxylon renaulti** Lignier.

1907 **Hamamelidoxylon renaulti** Lignier, p. 301, pls. XIX, XX, f. 44—52; XXI, f. 68; XXIII, f. 85, 93; text-fig. 3—5.

Remarks: Compared particularly with *Parrotia*. Two other fossil woods referred to the *Hamamelidaceae* have been described (see *Liquidambaroxylon*).

Occurrence: Tertiary; Antigua, Mexico.

Hauera Unger, 1845, p. 228.**Hauera americana** Unger.

1833 Dicotyledonous wood: Witham, p. 67, pl. XVI, f. 14.

1845 **Hauera americana** Unger, p. 228.

1850 **Hauera americana** Unger: Unger, p. 426.

1857a **Hauera americana** Unger: Unger, p. 17, pl. IV, f. 6, 7.

1882 **Hauera americana** Unger: Felix, p. 68.

1890 **Hauera americana** Unger: Kaiser, p. 30.

1894 **Hauera americana** Unger: Felix, p. 98.

Remarks: Referred by Unger to the *Aquilarineae*. Schenk (1890, p. 900) suggests a comparison with the *Caesalpineae*, and Felix (1894) with the *Aurantiaceae*.

Occurrence: Tertiary; Antigua, Mexico.

Hauera bornensis Engelhardt.

1870 **Hauera bornensis** Engelhardt, p. 49, pl. XV, f. 10—13.

Occurrence: Miocene; Saxony (Borna).

Hauera stiriaca Unger.

1845 **Hauera stiriaca** Unger, p. 229.

1850 **Hauera stiriaca** Unger: Unger, p. 426.

1857a **Hauera stiriaca** Unger: Unger, p. 17, pl. IV, f. 4, 5.

1882 **Hauera stiriaca** Unger: Felix, p. 68.

1890 **Hauera stiriaca** Unger: Kaiser, p. 30.

Occurrence: Miocene; Styria.

Helictoxylon Felix, 1882, p. 41.**Helictoxylon anomalum** Felix.

1883 **Helictoxylon anomalum** Felix, p. 66, pl. II, f. 4; III, f. 9.

1884 **Helictoxylon anomalum** Felix: Felix, p. 35.

1889 **Helictoxylon anomalum** Felix: Staub, p. 190.

1890 **Helictoxylon anomalum** Felix: Kaiser, p. 38.

Remarks: A liane of uncertain systematic position.

Occurrence: Tertiary; Hungary (Tapolcsan).

Helictoxylon luzonense Crié.

- 1889 **Helictoxylon luzonense** Crié, p. 87, pl. IX (XVII), f. 5—6.
Occurrence: Tertiary; Philippines (Luzon, Manilla, S. Juan del Monte).

Helictoxylon roemeri Felix.

- 1882 **Helictoxylon roemeri** Felix, p. 40.
1883a **Helictoxylon roemeri** Felix: Felix, p. 20, pl. III, f. 3, 7.
1883 **Helictoxylon roemeri** Felix: Hofmann, p. 92.
1890 **Helictoxylon roemeri** Felix: Kaiser, p. 38.
Remarks: Liane. Original in Munich Mus.
Occurrence: Probably Tertiary; Galicia (Tarnow), Mecklenburg ?

Helictoxylon schenki Felix.

- 1882 **Helictoxylon schenki** Felix, p. 62.
1883a **Helictoxylon schenki** Felix: Felix, p. 20, pl. III, f. 4, 5.
1890 **Helictoxylon schenki** Felix: Kaiser, p. 39.
1930 **Helictoxylon schenki** Felix: Schönfeld, p. 124.
Remarks: A silicified liane, which Schönfeld refers to the *Ternstroemiaceae*, and compares with *Ruyschioxylon* from Sumatra.
Occurrence: Tertiary; Java.

Helictoxylon speciosum Felix.

- 1882 **Helictoxylon speciosum** Felix, p. 66, pl. I, f. 1.
1883a **Helictoxylon speciosum** Felix: Felix, p. 18, pl. III, f. 2, 8.
1883 **Helictoxylon speciosum** Felix: Hofmann, p. 92.
Remarks: Silicified liane of uncertain affinity.
Occurrence: Tertiary; Antigua, ? Mecklenburg.

Helictoxylon tenerum Felix.

- 1882 **Helictoxylon tenerum** Felix, p. 67.
1883a **Helictoxylon tenerum** Felix: Felix, p. 19, pl. III, f. 1, 6.
1890 **Helictoxylon tenerum** Felix: Kaiser, p. 39.
Remarks: Silicified liane, affinities uncertain.
Occurrence: Tertiary; Antigua.

Helictoxylon wilcoxianum Berry.

- 1922 **Helictoxylon wilcoxianum** Berry, p. 18, pls. XVII, XVIII.
Occurrence: Eocene (Wilcox); Louisiana (Naborton).

Helictoxylon sp.

- 1882 **Helictoxylon** sp.: Hofmann, p. 92.
Remarks: Perhaps *H. speciosum* Felix.
Occurrence: Locality and horizon unknown. (Rostock Museum).

Hippocrateoxylon Hofmann, 1884, p. 179.

Hippocrateoxylon javanicum Hofmann.

- 1884 **Hippocrateoxylon javanicum** Hofmann, p. 179.
1890 **Hippocrateoxylon javanicum** Hofmann: Kaiser, p. 28.

1930 *Hippocrateoxylon javanicum* Hofmann: Schönfeld, p. 124.

Remarks: Silicified liane, compared with *Hippocratea scandens* and *H. viridis*. Original in Utrecht Univ. Coll.

Occurrence: Tertiary; Java (near Indramaju).

Hythia Stopes 1915, p. 277.

Hythia elgari Stopes.

1915 *Hythia elgari* Stopes, p. 278, pls. XXIX, XXX; text-fig. 85, 86.

Remarks: Affinities uncertain (? cf. *Fagus*). Originals in Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Lower Cretaceous (Aptian); England (Kent).

[**Jordania** Schenk, 1880.]

Remarks: Non *Jordania* Fiedler, Nova Acta, XXVI, 1857, p. 239; non *Jordania* Boissier 1849. See *Ebenoxylon ebenoides* (Schenk) and *E. tunetanum* (Fliche).

Juglandinium Unger, 1845, p. 241.

Remarks: The diagnosis was emended by Kraus (1882) who used the names *Juglandinium* and *Juglandoxylon* indifferently.

Juglandinium longiradiatum Vater.

1884 *Juglandinium longiradiatum* Vater, p. 841, pl. XXVIII, f. 15, 16.

1890 *Juglandinium longiradiatum* Vater: Kaiser, p. 14.

Occurrence: Lower Senonian; Germany (Harzburg).

Juglandinium mediterraneum Unger.

1845 *Juglandinium mediterraneum* Unger, p. 241.

1850 *Juglandinium mediterraneum* Unger: Unger, p. 472.

1852 *Juglandinium mediterraneum* Unger: Prokesch-Osten & Unger, p. 857.

1874 *Juglandinium mediterraneum* Unger: Schimper, p. 253.

1882 cf. *Juglandinium mediterraneum* Unger: Kraus, p. 5.

1882c cf. *Juglandinium mediterraneum* Unger: Kraus, p. 91.

1889 *Juglandinium mediterraneum* Unger: Staub, p. 189.

1890 *Juglandinium mediterraneum* Unger: Kaiser, p. 15.

1893 "*Juglandoxylon mediterraneum* Kraus": Meschinelli & Squinabol, p. 244.

1898 *Juglandinium mediterraneum* Unger: Fliche, p. 302.

Remarks: As Unger's wood was insufficiently described and not figured at all, Kraus hesitated to identify his Sicilian wood with it, but he did not make a new species. On p. 80 Kraus (1882c) refers to his wood as *Juglandoxylon*, on p. 87 as *Juglans*, and on p. 91 as "*Juglandinium* or *Juglandoxylon*". Nagel (1915, p. 65) includes *Mirbellites lesbius* Unger as a synonym of this species.

Occurrence: Tertiary; Mytilene, Hungary (Neugrad), ? Sicily (Sulphur beds of Girgenti).

Juglandinium schenki Felix.

1884 *Juglandinium schenki* Felix, p. 30, pl. II, f. 1—3.

1889 *Juglandinium schenki* Felix: Staub, p. 189.

1890 *Juglandinium schenki* Felix: Kaiser, p. 15.

Remarks: Original in Min. Mus. Leipzig.

Occurrence: Tertiary; Hungary.

***Juglandinium triebeli* (Caspary) Nagel.**

- 1888 *Juglans triebelii* Caspary, p. 42.
1889 *Juglans triebelii* Caspary: Caspary, p. 50, pl. IX, f. 12, 13; X, f. 1—9.
1890 *Juglans triebelii* Caspary: Kaiser, p. 15.
1915 *Juglandinium triebelii* (Caspary) Nagel, p. 65.
Remarks: Compared with *Pterocarya caucasica*. Originals in Mus. Bot. Gart., Königsberg. Schuster (1908, p. 150) compares it with *Juglans australis*.
Occurrence: ? Tertiary; Prussia (? Elbing).

***Juglandinium wichmanni* (Hofmann).**

- 1884 *Juglandoxylon wichmanni* Hofmann, p. 187 (36).
1890 *Juglandoxylon wichmanni* Hofmann: Kaiser, p. 15.
Occurrence: Locality and horizon unknown. (Utrecht Univ. Coll.).

***Juglandinium zuriense* (Falqui).**

- 1906 *Juglansoxylon zuriensis* Falqui, p. 26, pl. I, f. 2.
Occurrence: Miocene; Sardinia.

***Juglandinium* sp.**

- 1884 *Juglandinium* sp.: Vater, p. 842.
1890 *Juglandinium* sp.: Kaiser, p. 15.
Occurrence: Lower Senonian; Brunswick (Helmstedt).

? *Juglandinium* sp.

- 1888 *Carya*: Dawson, p. 33.
Remarks: No figures nor description. Requires confirmation.
Occurrence: Upper Cretaceous (Fort Pierre Series); Western Canada (Head of Swift Current).

***Juglandinium* ? sp.**

- 1930a *Juglandinium*?: Schönfeld, p. 70.
Occurrence: Oligocene; Saxony (Deutzen).

[***Juglandoxylon* Kraus, 1882c, p. 80.]**

Synonym of ***Juglandinium***.

[***Juglans triebelii* Caspary.]**

Synonym of ***Juglandinium triebelii***, q. v.

[***Juglansoxylon* Falqui, 1906.]**

Synonym of ***Juglandinium***, q. v. for ***J. zuriense***.

***Jugloxylon* Stopes & Fujii, 1910, p. 62.**

***Jugloxylon hamoanum* Stopes & Fujii.**

- 1910 *Jugloxylon hamoanum* Stopes & Fujii, p. 62, pl. VII, f. 48.
Remarks: Affinities uncertain. Does not appear to resemble *Juglans* very closely, in spite of the suggestion embodied in the name.
Occurrence: Upper Cretaceous; Japan (Hokkaido).

[**Junghuhnites** Goeppert, 1854, p. 54.][**Junghuhnites javanicus** Goeppert.]1854 **Junghuhnites javanicus** Goeppert, p. 54, pl. II, f. 11—16.1890 **Junghuhnites javanicus** Goeppert: Kaiser, p. 35.

Remarks: Kräusel (1925, p. 333) suggested that this wood might be Dipterocarpaceous, but later (1926, p. 1) decided that it was quite unidentifiable. The originals appear to be lost.

Occurrence: Tertiary; Java.

Klippsteinia Unger, 1845, p. 234.**Klippsteinia medullaris** Unger.1845 **Klippsteinia medullaris** Unger, p. 234.1850 **Klippsteinia medullaris** Unger: Unger, p. 448.1857 **Klippsteinia medullaris** Unger: Unger, p. 12, pl. III, f. 8—10.1890 **Klippsteinia medullaris** Unger: Kaiser, p. 25.1894 **Klippsteinia medullaris** Unger: Felix, p. 98.

Remarks: Referred by Unger to the *Aurantiaceae*. Schenk (1890, p. 899) remarks that a similar structure is also to be found in the *Celastraceae*. Felix (1894) doubts the reference to the *Aurantiaceae*.

Occurrence: Tertiary; Austria (Thal near Graz).

[**Kloedenia** Goeppert, 1839.][**Kloedenia quercoides** Goeppert.]

Synonym of **Quercinium quercoides**, q. v.

Laurinium Unger, 1845, p. 227.

Remarks: Felix (1886, p. 490 and 1887a, p. 156) attempted to divide Unger's genus into two: (a) *Laurinium* s. str., in which there are no secretory canals in the rays, and (b) *Perseoxyton*, secretory canals present in the rays. The latter included species which Felix had previously named *Laurinoxylon* (1883). Vater (1884) reverted to *Laurinium* and emended Unger's diagnosis. Knoblauch, followed by Schuster and Gothan, pointed out that the presence of secretory canals was by no means confined to *Persea* among recent *Lauraceae*, and Knoblauch (quoted by Schuster) concluded that lauraceous genera could not be distinguished by their wood anatomy. Nevertheless Schuster (1908, 1909) endeavoured to distinguish certain woods under the name *Ocoteoxyton*, but Gothan does not consider that this is justifiable.

Gothan (followed by Kräusel) rejects the name *Laurinium* in favour of *Laurinoxylon*, on what seem to me inadequate grounds.

Laurinium albiense (Fliche).1905 **Laurinoxylon albiense** Fliche, p. 356, pl. X, f. 2, 3.

Occurrence: Albian; Madagascar.

Laurinium algovicum (Schuster).1909 **Ocoteoxyton algovicum** Schuster, p. 55.

Remarks: Compared with the recent genus *Ocotea*, but Gothan (1908, p. 18) doubts whether the separation of *Ocoteoxyton* from *Laurinium* is justifiable. Cf. *Laurinium (Ocoteoxyton) tigurinum*.

Occurrence: Upper Oligocene; Bavaria (Algäu).

Laurinium brunswicense Vater.

- 1884 *Laurinium brunswicense* Vater, p. 845, pl. XXIX, f. 22—24.
 1887a *Laurinium brunswicense* Vater: Felix, p. 157.
 1890 *Laurinium brunswicense* Vater: Kaiser, p. 21.
 1908 *Laurinium brunswicense* Vater: Schuster, p. 144.

Remarks: Compared with *Laurus nobilis*.

Occurrence: Lower Senonian (derived from); Brunswick (Helmstedt).

Laurinium californicum (Platen).

- 1908 *Perseoxylon californicum* Platen, p. 52.

Occurrence: Tertiary; California.

Laurinium desioi (Chiarugi).

- 1929 *Laurinoxylon desioi* Chiarugi, p. 423, pl. XLVII, f. 1—3.

- 1929a *Laurinoxylon desioi* Chiarugi: Chiarugi, p. 558.

Occurrence: Miocene; Libyan desert (Giarabub), Sirtica.

Laurinium diluviale (Unger).

- 1842 *Ulmium diluviale* Unger, p. 101.
 1842a *Ulmium diluviale* Unger: Unger, p. 174.
 1845 *Ulmium diluviale* Unger: Unger, p. 219.
 1847 *Ulmium diluviale* Unger: Unger, p. 97, pl. XXV, f. 6—9.
 1850 *Ulmium diluviale* Unger: Unger, p. 412.
 1868 *Ulmium diluviale* Unger: Cramer, p. 175.
 1872 *Ulmium diluviale* Unger: Schimper, p. 724.
 1882 *Betulium diluviale* (Unger) Felix, p. 40.
 1883 *Laurinoxylon diluviale* (Unger) Felix, p. 59, pl. II, f. 1, 3; III, f. 1.
 1884 *Laurinoxylon diluviale* (Unger): Felix, p. 28.
 1886 *Perseoxylon diluviale* (Unger) Felix, p. 490.
 1887a *Perseoxylon diluviale* (Unger): Felix, p. 157.
 1890 *Perseoxylon diluviale* (Unger): Kaiser, p. 23.
 1890 *Betuloxylon diluviale* (Unger) Lakowitz, p. 29.

Remarks: Conwentz (1868) and Kaiser (1879) had already expressed doubts as to the correctness of Unger's ascription of this wood, and Felix at first, following Cramer's suggestion, renamed it *Betulium*, but after an examination of some of the original material he identified it as lauraceous. (See also remarks on *Betulium* sp. from Cracow).

In spite of Felix's work, Lakowitz (1890) refers to it as *Betuloxylon diluviale*, and then states that he does not consider it to be a birch wood!

Felix stated that the wood closely resembled that of the recent *Persea gratissima* Gaertn., and later he included it in his genus *Perseoxylon*. Gothan and others doubt the possibility of distinguishing different genera of the *Lauraceae* by the characters of the wood, a view which has been adopted here, so that this species becomes *Laurinium diluviale* (Unger).

Occurrence: Tertiary; Bohemia (Joachimsthal).

Laurinium eberi (Platen).

- 1908 *Perseoxylon eberi* Platen, p. 135.

Occurrence: Tertiary (Miocene?); Colorado (Bijou Basin).

Laurinium guatemalense Unger.

- 1850 *Laurinium guatemalense* Unger, p. 425.
 1887a *Laurinium guatemalense* Unger: Felix, p. 157.
 1890 *Laurinium guatemalense* Unger: Kaiser, p. 21.
 1908 *Laurinium guatemalense* Unger: Schuster, p. 143.
 Remarks: Included in *Laurinium* with a query by Felix.
 Occurrence: ? Tertiary; Guatemala (Rio Payres).

Laurinium haasi Wetzel.

- 1913 *Laurinium haasi* Wetzel, p. 21, pl. I.
 Occurrence: ? Upper Cretaceous; Germany (Holtenau).

Laurinium lesquerianum (Knowlton).

- 1891 *Laurinoxylon lesqueriana* Knowlton, p. 258, pl. X, f. 3, 4; XI, f. 3, 4.
 1924 *Laurinoxylon lesqueriana* Knowlton: Berry, p. 85.
 Remarks: Originals in U. S. Nat. Mus.
 Occurrence: Eocene (Yegua); Arkansas.

Laurinium meyeri Felix.

- 1886 *Laurinium meyeri* Felix, p. 488, pl. XII, f. 4, 7, 8.
 1887a *Laurinium meyeri* Felix: Felix, p. 157.
 1890 *Laurinium meyeri* Felix: Kaiser, p. 21.
 1908 *Laurinium meyeri* Felix: Schuster, p. 143.
 Remarks: Originals in Min. Mus. Dresden. Recorded as lauraceous wood, on the identification of Conwentz, by Frenzel (Min. Mitt., 1877, p. 308).
 Occurrence: Tertiary; New Guinea (Astrolabe Bay).

Laurinium nectandrioides (Kräusel & Schönfeld).

- 1924 *Laurinoxylon nectandrioides* Kräusel & Schönfeld, p. 272, pl. XXII, f. 19—21; text-fig. 9—17.
 Remarks: Agrees closely with the wood of *Nectandra*. Probably the same as the wood referred to by Schönfeld (1925, p. 19) as cf. *Nectandra*.
 Occurrence: Miocene; Holland (S. Limburg).

Laurinium perseoides (Caspary).

- 1888 *Laurus perseoides* Caspary, p. 43.
 1889 *Laurus perseoides* Caspary: Caspary, p. 67, pl. XII, f. 6—11; XIII, f. 1—5.
 1890 *Laurus perseoides* Caspary: Kaiser, p. 21.
 Remarks: Compared with *Persea gratissima*. Original in Mus. Bot. Gart., Königsberg.
 Occurrence: Tertiary; Prussia (Palmnicken).

Laurinium primigenium (Schenk) Felix.

- 1883 *Laurinoxylon primigenium* Schenk, p. 11, pl. III, f. 10; V, f. 15, 16.
 1884 *Laurinoxylon primigenium* Schenk: Felix, p. 28.
 1887a *Laurinium primigenium* (Schenk) Felix, p. 157.
 1890 *Laurinoxylon primigenium* Schenk: Kaiser, p. 22.
 1908 *Laurinoxylon primigenium* Schenk: Schuster, p. 143.



Remarks: Kaiser suggests a similarity to the wood of the *Rubiacae*. Schuster doubts whether it is lauraceous.

Occurrence: Oligocene ?; Egypt (near Cairo).

Laurinium pulchrum (Knowlton).

1899 *Laurinoxylon pulchrum* Knowlton, p. 765, pl. CXVI, CXIX, f. 3—5, CXX, f. 1.

1908 *Laurinoxylon pulchrum* Knowlton: Platen, p. 127.

Occurrence: Miocene; Yellowstone National Park.

Laurinium radiatum Schönfeld.

1919 *Laurinium radiatum* Schönfeld, p. 3, pl. I, f. 1—6.

Remarks: Compared with *L. tigurinum* (Schuster), and also with *Carpinoxylon pfefferi* Platen, which, it is suggested, may possibly be lauraceous.

Occurrence: Tertiary; Dresden.

Laurinium tigurinum (Schuster).

1908 *Ocoteoxylon tigurinum* Schuster, p. 139, pl. II.

1919 *Ocoteoxylon tigurinum* Schuster: Schönfeld, p. 8.

Remarks: Gothan (1908, p. 18) does not consider that the separation of *Ocoteoxylon* from *Laurinium* is justified. It seems better to use the more comprehensive form-genus, though this species may, as Schuster claims, show points of contact with the recent *Ocotea*. Among fossils it is near *L. biseriatum*.

Occurrence: Eocene (Flysch); Tegernsee.

Laurinium triseriatum (Caspary).

1888 *Laurus triseriata* Caspary, p. 43.

1889 *Laurus triseriata* Caspary: Caspary, p. 60, pl. XI, f. 6—12; XII, f. 1—5.

1890 *Laurus triseriata* Caspary: Kaiser, p. 22.

Remarks: Compared with *Laurus nobilis* and *Mithridatea erecta*. Original in Mus. Bot. Gart., Königsberg.

Occurrence: Tertiary; Prussia.

Laurinium uniseriatum (Gothan).

1908 *Laurinoxylon uniseriatum* Gothan, p. 16, pl. II, f. 3—11.

1924 *Laurinoxylon "uniradiatum"* Gothan: Kräusel, p. 25, pl. IV, f. 1—4.

Remarks: Kräusel's specific name is presumably a mistake for *L. uniseriatum*.

Occurrence: Upper Cretaceous or Tertiary; Antarctica (Seymour Island).

Tertiary; Patagonia.

Laurinium wilcoxianum (Berry).

1922 *Laurinoxylon wilcoxianum* Berry, p. 19, pl. XIII.

Remarks: The statement that this species was present in the Yegua of Texas probably refers to the later described *L. bakeri* Berry.

Occurrence: Eocene (Wilcox); Louisiana (Naborton), Missouri (Dabnez), Tennessee (Bolivar).

Laurinium xyloides Unger.

- 1845 *Laurinium xyloides* Unger, p. 228.
 1850 *Laurinium xyloides* Unger: Unger, p. 425.
 1887a *Laurinium xyloides* Unger: Felix, p. 157.
 1890 *Laurinium xyloides* Unger: Kaiser, p. 22.
 1893 *Lauroxylon xyloides* (Unger) Meschinelli & Squinabol, p. 303.
 1908 *Laurinium xyloides* Unger: Schuster, p. 143.

Remarks: Schuster states that this species is insufficiently described. Felix includes it in *Laurinium* with a query.

Occurrence: Pliocene: Italy (Vicentino).

Laurinium sp.

- 1857 *Laurus* sp.: Meneghini, pp. 439, 548.
 1893 *Lauroxylon* sp.: Meschinelli & Squinabol, p. 303.

Remarks: Compared with *Laurus benzoin*.

Occurrence: Tertiary; Sardinia.

Laurinium sp.

- 1891 *Laurinoxylon*?: Knowlton, p. 259.
 1924 *Laurinoxylon* ? sp. Knowlton: Berry, p. 188.
 Remarks: Very poorly preserved. Original in U. S. Nat. Mus.
 Occurrence: Eocene; Arkansas (Red Bluff).

Laurinium ? sp.

- 1908 *Laurinoxylon* ? sp.: Gothan, p. 19, pl. II, f. 12, 13.
 Occurrence: ? Upper Cretaceous or Tertiary; Antarctica (Seymour Island).

Laurinium sp.

- 1930a *Laurinium*: Schönfeld, p. 70.
 1930a *Perseoxylon*: Schönfeld, p. 70.
 Occurrence: Oligocene; Saxony (Ragewitz, Flöz, Quatitz).

[Laurinoxylon Felix, 1883, p. 59.]

Synonym of *Laurinium*, q. v. for all species described under *Laurinoxylon*.

[Lauroxylon Schenk, 1890, p. 899.]

Synonym of *Laurinium*, q. v.

Lillia Unger, 1842, p. 102.**Lillia viticulosa** Unger.

- 1842 *Lillia viticulosa* Unger, p. 102.
 1842a *Lillia viticulosa* Unger: Unger, p. 178.
 1845 *Lillia viticulosa* Unger: Unger, p. 263.
 1845 *Lillia viticulosa* Unger: Corda, p. 49, pl. LX, f. 1—3.
 1850 *Lillia viticulosa* Unger: Unger, p. 477.
 1883 *Lillia viticulosa* Unger: Felix, p. 64.
 1884 *Lillia viticulosa* Unger: Felix, p. 33, pl. IV, f. 5, 6.
 1889 *Lillia viticulosa* Unger: Staub, p. 189.

1890 *Lillia viticulosa* Unger: Kaiser, p. 23.

1911 *Lillia viticulosa* Unger: Schuster, p. 540.

1930 *Lillia viticulosa* Unger: Schönfeld, p. 124.

Remarks: Corda suggested an affinity with the *Zygophyllaceae*, but Felix pointed out the close resemblance of this liane to *Coscintium fenestratum* and put it in the *Menispermaceae*. Originals in Nat. Mus. Vienna and Geol. Mus. Dresden.

Occurrence: Tertiary; Hungary (Rauca, Gyepüfűzes).

Liquidambaroxylon Felix, 1884, p. 24.

Liquidambaroxylon lecointreae (Houlbert).

1910 *Ambaroxylon lecointreae* Houlbert, p. 74, pl. VII, f. 17—20.

Remarks: Closely compared with *Liquidambar*, and therefore falling within Felix's form-genus.

Occurrence: Middle Miocene; Touraine.

Liquidambaroxylon speciosum Felix.

1884 *Liquidambaroxylon speciosum* Felix, p. 24, pl. III, f. 2—4; IV, f. 4.

1889 *Liquidambaroxylon speciosum* Felix: Staub, p. 189.

1890 *Liquidambaroxylon speciosum* Felix: Kaiser, p. 29.

Remarks: Felix puts this species very close to *Liquidambar styraciflua*. Schenk (1890, p. 903) says "Bei *Liquidambaroxylon* vermisse ich die in dem primären Holze vorkommenden Secretgänge". Originals in Mus. Ungar. Geol. Reichsanst.

Occurrence: Tertiary; Hungary (Medgyaszó).

Magnolia L.

[*Magnolia laxa* Caspary.]

See *Dryoxylon laxum*.

[*Magnolites Tuzson, 1909.*]

[*Magnolites silvatica* Tuzson.]

See *Dryoxylon silvaticum*.

Meyenites Unger, 1842, p. 102.

Meyenites aequimontanus Unger.

1842 *Meyenites aequimontanus* Unger, p. 102.

1842a *Meyenites aequimontanus* Unger: Unger, p. 177.

1845 *Meyenites aequimontanus* Unger: Unger, p. 261.

1850 *Meyenites aequimontanus* Unger: Unger, p. 522.

1854 *Meyenites aequimontanus* Unger: Unger, p. 183, pl. VII, f. 4—6.

1890 *Meyenites aequimontanus* Unger: Kaiser, p. 35.

Occurrence: Miocene; Styria (Gleichenberg).

[*Miquelites* Goepfert.]

[*Miquelites elegans* Goepfert.]

1854 *Miquelites elegans* Goepfert, p. 56, pl. I, f. 6, 7, 7a.

1890 *Miquelites elegans* Goepfert: Kaiser, p. 35.

Remarks: Kräusel states (1926) that although this wood shows some resemblance to that of the *Dipterocarpaceae*, it must be regarded as quite unidentifiable.

Occurrence: Tertiary; Java.

Mirbellites Unger, 1845, p. 241.

Mirbellites lesbius Unger.

- 1845 *Mirbellites lesbius* Unger, p. 242.
 1850 *Mirbellites lesbius* Unger: Unger, p. 472.
 1852 *Mirbellites lesbius* Unger: Prokesch-Osten & Unger, p. 857.
 1890 *Mirbellites lesbius* Unger: Kaiser, p. 15.
 1898 *Mirbellites lesbius* Unger: Fliche, p. 302.

Remarks: Referred by Unger to the *Juglandaceae*. Nagel (1915, p. 65) includes it in the synonymy of *Juglandinium mediterraneum*, but there is no evidence in favour of this.

Occurrence: Tertiary; Lesbos.

Mirbellites schuchii Unger.

- 1850 *Mirbellites schuchii* Unger, p. 473.
 1890 *Mirbellites schuchii* Unger: Kaiser, p. 16.
 Occurrence: Horizon and locality unknown.

Mohlites Unger, 1842, p. 101.

Mohlites cribrosus Unger.

- 1842 *Mohlites cribrosus* Unger, p. 101.
 1842a *Mohlites cribrosus* Unger: Unger, p. 176.
 1845 *Mohlites cribrosus* Unger: Unger, p. 265.
 1850 *Mohlites cribrosus* Unger: Unger, p. 525.
 1889 *Mohlites cribrosus* Unger: Staub, p. 191.
 1890 *Mohlites cribrosus* Unger: Kaiser, p. 32.
 Occurrence: Tertiary; Hungary (Libethen).

Mohlites parenchymatosus Unger.

- 1842 *Mohlites parenchymatosus* Unger, p. 101.
 1842a *Mohlites parenchymatosus* Unger: Unger, p. 176.
 1845 *Mohlites parenchymatosus* Unger: Unger, p. 265.
 1847 *Mohlites parenchymatosus* Unger: Unger, p. 3, pl. I, f. 3a.
 1850 *Mohlites parenchymatosus* Unger: Unger, p. 525.
 1854 *Mohlites parenchymatosus* Unger: Unger, p. 182, pl. VI, f. 14—16.
 1890 *Mohlites parenchymatosus* Unger: Kaiser, p. 32.

Remarks: Referred by Unger to the *Leguminosae*. The figure in Chloris Protogaea (1847) is in illustration of a fungus (*Nyctomyces antediluvianus*) with which the wood was infested.

Occurrence: Miocene; Styria (Gleichenberg).

Nectandra Roland.

See *Laurinium nectandrioides*.

Nicolia Unger, 1842, p. 102.

Nicolia aegyptiaca Unger.

- 1842 *Nicolia aegyptiaca* Unger, p. 102.
 1842a *Nicolia aegyptiaca* Unger: Unger, p. 177.
 1845 *Nicolia aegyptiaca* Unger: Unger, p. 262.

- 1847 (Fossil wood): Unger, pl. I, f. 7.
 1850 *Nicolia aegyptiaca* Unger: Unger, p. 523.
 1859 *Nicolia aegyptiaca* Unger: Unger, p. 213, pl. I, f. 1, 2.
 1866 *Nicolia aegyptiaca* Unger: Unger, p. 289, pl. I, f. 1—7.
 1870 *Nicolia aegyptiaca* Unger: Carruthers, p. 307, pl. XIV, f. 3, 4.
 1880 *Nicolia aegyptiaca* Unger: Schenk, p. 657.
 1883 *Nicolia aegyptiaca* Unger: Schenk, p. 8, pl. III, f. 7—9; IV, f. 11.
 1884a *Nicolia wiedemanni* Hofmann, p. 485, pl. III, f. 1.
 1885 *Nicolia aegyptiaca* Unger: Quenstedt, p. 1175, pl. C, f. 49.
 1888 *Nicolia aegyptiaca* Unger: Schenk, pp. 19, 23.
 1890 *Nicolia aegyptiaca* Unger: Kaiser, p. 31.
 1890 *Nicolia wiedemanni* Hofmann: Kaiser, p. 36.
 1890 *Nicolia aegyptiaca* Unger: Lange, p. 673.
 1893 *Nicolia aegyptiaca* Unger: Thomas, pp. 3, 4.
 1901 *Nicolia aegyptiaca* Unger: Blanckenhorn, pp. 98, 102.
 1910 *Nicolia aegyptiaca* Unger: Schuster, pp. 5—8, pl. II, f. 9—11; III, f. 13—15.
 1925 *Nicolia aegyptiaca* Unger: Edwards, p. 171.
 1929 *Nicolia aegyptiaca* Unger: Chiarugi, p. 410, pl. XLV, f. 5; XLVI, f. 1, 3, 5; text-fig. 38, 39, 41, 46.

Remarks: For discussions on the modes of occurrence and fossilization as well as the distribution of *Nicolia* and other woods of the so-called "petrified forests" of Egypt, see, in addition to papers above: Schweinfurth (1882), Kuntze (1895), Barron (1905) and Stromer (1924).

Nicol (1835) mentions dicotyledonous wood from Egypt, some of which, as is shown from an examination of his original specimens in the British Museum (Nat. Hist.) can be identified as *Nicolia aegyptiaca*.

Unger (1847) figured a fungus (*Nyctomyces entoxylinus*) in a dicotyledonous wood from Asserac, Egypt, which he afterwards stated was *N. aegyptiaca* (1859, p. 221).

The figures given by Quenstedt (1885) of a specimen from near Cairo are poor and unrecognizable.

The record by Lange from the Lower Senonian of Aix-la-Chapelle (no figures) requires confirmation.

Schuster includes *N. wiedemanni* Hofmann in *N. aegyptiaca*, and Chiarugi follows this.

Schuster considers that pitted elements figured by Schacht (Ann. Sci. Nat. Bot., XIII, 1860) as "*lignum Leguminosarum fossilium*" (no locality stated) must have belonged to *N. aegyptiaca*.

Occurrence: (See map given by Chiarugi, 1929).

Upper Cretaceous (Nubian Sandstone); Libyan Desert (near Regenfeld).

? Lower Senonian; ? Aix-la-Chapelle.

Eocene (Daban Beds); British Somaliland.

Oligocene; Egypt (near Cairo and various other localities).

Miocene; Libyan Desert (Moghara; Wadi Faregh; between Fayum and Baharia; Giarabub).

Pliocene; Egypt (Wadi Natrun, Garet Muluk) [Stromer (1924, p. 20) thinks this wood may have been derived from the Lower Miocene].

Tunisia (Ain Cherichira and Bled-Douara).

Tertiary (horizon uncertain); Abyssinia (Wadla, Woro, Haimano); Italian Somaliland, West coast of Africa near Huissi Aissa.

Nicolia caledonica Crié.

- 1889 *Nicolia caledonica* Crié, p. 81, pl. V (XIII), f. 1, 2; VI (XIV), f. 1—8.

Occurrence: ? Pliocene; New Caledonia (Ducos I.).

[*Nicolia minor* Hofmann.]

Synonym of *Caesalpinioxylon oweni* (Carr.), q. v.

Nicolia giarabubensis Chiarugi.

1929 *Nicolia giarabubensis* Chiarugi, p. 418, pl. XLV, f. 6; XLVI, f. 2, 4, 6.

Occurrence: Miocene; Libyan Desert (Giarabub).

Nicolia moresneti Hovelacque.

1888 *Achenosaurus multidens* Smets, pars. [Not seen].

1890 *Nicolia moresneti* Hovelacque, p. 63, pl. III, f. 2; text-fig. 2-8.

Remarks: See remarks on *Achenoxylon multidens*. Compared by Hovelacque with the *Piperaceae*.

Occurrence: Upper Cretaceous; Belgium (Moresnet).

[*Nicolia oweni* Carruthers.]

Synonym of *Caesalpinioxylon oweni*, q. v.

[*Nicolia tunetana* Crié.]

1889 *Nicolia tunetana* Crié, p. 81 (footnote).

Remarks: Nomen nudum.

Occurrence: Pliocene; Tunis.

[*Nicolia wiedemanni* Hofmann.]

Synonym of *N. aegyptiaca*, q. v.

Nicolia zelandica Unger.

1864 *Nicolia zelandica* Unger, p. 13, pl. V, f. 2a-b.

Remarks: Schenk (1890, p. 901) doubts whether the generic attribution of this wood is correct.

Occurrence: Tertiary; New Zealand (Hunua Range).

Nicolia ? sp.

1888 *Nicolia* ? : Fliche, p. 571.

Remarks: Cf. the record of *N. aegyptiaca* in Thomas (1893).

Occurrence: Tertiary; Tunisia (Ain Cherichira).

Nothofagoxylon Gothan, 1908.

Nothofagoxylon scalariforme Gothan.

1908 *Nothofagoxylon scalariforme* Gothan, p. 20, pl. II, f. 14-18.

1924 *Nothofagoxylon scalariforme* Gothan: Kräusel, p. 19, pl. III, f. 1-3.

Remarks: Gothan compares this species particularly with the living *Nothofagus betuloides* Bl.

Occurrence: ? Upper Cretaceous or Tertiary; Antarctica (Seymour Island) and Patagonia.

Nothofagoxylon sp.

1924 **Nothofagoxylon** sp. (*scalariforme* ?): Kräusel, p. 23, pl. I, f. 13; III, f. 4—9.

1924 **Nothofagoxylon** sp.: Kräusel, p. 29.

Occurrence: Tertiary; Chiloe and Tierra del Fuego.

[**Ocoteoxylon** Schuster, 1908, p. 139.]

Included in **Laurinium**, q. v.

Ornoxylon Felix, 1882, p. 35.

Ornoxylon fraxinoides Felix.

1882 **Ornoxylon fraxinoides** Felix, p. 35.

1890 **Ornoxylon fraxinoides** Felix: Kaiser, p. 34.

Remarks: Silicified wood belonging to the *Oleaceae*, resembling the wood of *Fraxinus*. Original in Munich Museum.

Occurrence: Tertiary; near Cracow.

Palackya Crié, 1889, p. 87.

Palackya philippinense Crié.

1889 **Palackya philippinense** Crié, p. 87, pl. IX, (XVII), f. 1, 2.

1890 **Palackya philippinense** Crié: Schenk, p. 904.

Remarks: According to Schenk the structure suggests that of several *Caesalpiniaceae*.

Occurrence: Mio-Pliocene; Philippines (Manilla, S. Juan del Monte).

Paraphyllanthoxylon Bailey, 1924, p. 446.

Paraphyllanthoxylon arizonense Bailey.

1924 **Paraphyllanthoxylon arizonense** Bailey, p. 446, pl. XV.

Remarks: Referred doubtfully to the *Euphorbiaceae*, and compared with *Phyllanthus* and *Bridelia*. Might well have been included in *Euphorbioxylon* Felix.

Occurrence: Cretaceous (Colorado Group); Arizona.

Parinarioxylon Pfeiffer & van Heurn, 1928, p. 1011.

Parinarioxylon itersonii Pfeiffer & van Heurn.

1928 **Parinarioxylon itersonii** Pfeiffer & van Heurn, p. 1011, pl. II, f. 2, 3.

Remarks: Compared with *Parinarium sumatranum* (*Rosaceae*, *Chrysobalanoideae*).

Occurrence: Tertiary; Java (Bolang).

Pataloxylon Sahni, 1920, p. 29.

Pataloxylon porosum Sahni.

1920 **Pataloxylon porosum** Sahni, p. 31, pl. III, f. 12—15.

Remarks: Affinities uncertain. Type in Queensland Geol. Surv. Coll.

Occurrence: Tertiary; Queensland (Wolston near Brisbane).

Pataloxylon scalariforme Sahn.

1920 **Pataloxylon scalariforme** Sahn, p. 29, pl. I, f. 6; II, f. 8—11; text-fig. 10.

Remarks: Affinities uncertain. Type in Queensland Geol. Surv. Coll.

Occurrence: Tertiary; Queensland (Mt. Meerschaum).

[**Perseoxylon** Felix, 1886, p. 490.]

Included in **Laurinium**, q. v.

Petzholdia Unger, 1842, p. 102.

Petzholdia major Unger.

1845 **Petzholdia major** Unger, p. 260.

1850 **Petzholdia major** Unger: Unger, p. 521.

1890 **Petzholdia major** Unger: Kaiser, p. 36.

Occurrence: Tertiary; West Indies.

Petzholdia polonica Unger.

1850 **Petzholdia polonica** Unger, p. 522.

1890 **Petzholdia polonica** Unger: Kaiser, p. 36.

Occurrence: Miocene; Galicia (Viniki).

Petzholdia tropica Unger.

1833 **Dicotyledonous** wood: Witham, p. 67, pl. XVI, f. 12, 13.

1842 **Petzholdia tropica** Unger, p. 102.

1842a **Petzholdia tropica** Unger: Unger, p. 176.

1845 **Petzholdia tropica** Unger: Unger, p. 260.

1850 **Petzholdia tropica** Unger: Unger, p. 521.

1890 **Petzholdia tropica** Unger: Kaiser, p. 36.

Occurrence: Tertiary; Antigua.

[**Phegonium** Unger, 1842.]

Remarks: The name *Fegonium* was first spelt *Phegonium*.

Piccolominites Unger, 1845, p. 262.

Piccolominites sardus Unger.

1845 **Piccolominites sardus** Unger, p. 262.

1850 **Piccolominites sardus** Unger: Unger, p. 523.

1890 **Piccolominites sardus** Unger: Kaiser, p. 36.

Remarks: Stated to resemble *Nicolia*.

Occurrence: Tertiary; Sardinia.

Piper L.

cf. **Piper** sp.

1844 cf. **Piper** sp.: Bowerbank, p. 16.

Remarks: Pitted vessels and other elements regarded as belonging to *Piper*.

Occurrence: Eocene (London Clay); England.

Piperites Goeppert.**Piperites hasskarlianus** Goeppert.

1854 **Piperites hasskarlianus** Goeppert, p. 40, pl. III, f. 20—23.

1890 **Piperites hasskarlianus** Goeppert: Kaiser, p. 17.

Remarks: Kräusel (1925, p. 332, 1926, p. 1) regards this wood as quite unidentifiable, and states that the originals seem to be lost. The generic name was also used for leaf impressions.

Occurrence: Tertiary; Java.

Plataninium Unger, 1842, p. 101.

em. Vater, 1884, p. 842.

Plataninium acerinum Unger.

1842 **Plataninium acerinum** Unger, p. 101.

1842a **Plataninium acerinum** Unger: Unger, p. 174.

1845 **Plataninium acerinum** Unger: Unger, p. 222.

1847 **Plataninium acerinum** Unger: Unger, p. 133, pl. XLVII, f. 8—10.

1850 **Plataninium acerinum** Unger: Unger, p. 415.

1886 ? **Plataninium aceroides** (Goeppert) Windisch, p. 20.

1890 **Plataninium acerinum** Unger: Kaiser, p. 18.

Remarks: The wood recorded by Windisch from the Tertiary of Iceland is said to be similar to Unger's species, and also to the wood described by Schroeter as *Platanus aceroides* (see *Plataninium* sp.).

Occurrence: Formation and locality unknown.

Plataninium boreale (Caspary).

1888 **Platanus borealis** Caspary, p. 42.

1889 **Platanus borealis** Caspary: Caspary, p. 47, pl. IX, f. 1—11.

1890 **Platanus borealis** Caspary: Kaiser, p. 19.

Remarks: Stated to be nearer than *P. klebsii* to the living species of *Platanus*. Originals in Mus. Bot. Gart., Königsberg.

Occurrence: Tertiary; West Prussia (Plietnitz).

Plataninium crystallophilum Platen.

1908 **Plataninium crystallophilum** Platen, p. 111.

Occurrence: Tertiary; Arizona.

Plataninium haydeni Felix.

1896 **Plataninium haydeni** Felix, p. 251.

1899 **Plataninium haydeni** Felix: Knowlton, p. 767, pl. CXX, f. 3—5.

1908 **Plataninium haydeni** Felix: Platen, p. 129.

Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

Plataninium klebsii (Caspary).

1888 **Platanus klebsii** Caspary, p. 41.

1889 **Platanus klebsii** Caspary: Caspary, p. 40, pl. VII, f. 11, 12; VIII, f. 1—24.

1890 **Platanus klebsii** Caspary: Kaiser, p. 19.

Remarks: Compared with *Platanus occidentalis*. Original in Mus. Bot. Gart., Königsberg.

Occurrence: Oligocene; Prussia (Palmnicken).

Plataninium knowltoni Platen.1908 *Plataninium knowltoni* Platen, p. 130.

Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

Plataninium megapolitanum (Hofmann) Vater.1883 *Fegonium megapolitanum* Hofmann, p. 88.1884 *Plataninium megapolitanum* (Hofmann) Vater, p. 838.1887a *Plataninium megapolitanum* (Hofmann): Felix, p. 148.1890 *Plataninium megapolitanum* (Hofmann): Kaiser, p. 19.

Occurrence: Tertiary ?; Mecklenburg.

Plataninium pacificum Platen.1908 *Plataninium pacificum* Platen, p. 65.

Occurrence: Miocene; California (Nevada County).

Plataninium porosum Felix.1887 *Plataninium porosum* Felix, p. 146, pl. XXVII, f. 6.1889 *Plataninium porosum* Felix: Staub, p. 188.1890 *Plataninium porosum* Felix: Kaiser, p. 19.1894 *Plataninium porosum* Felix: Felix, p. 101.

Remarks: Root-wood. Originals in Mus. Ungar. Geol. Reichsanst.

Occurrence: Eocene (Sungait series); Caucasus (Apscheron).

? Tertiary; Hungary (Nagy-Almas, Hunyad).

Plataninium regulare Felix.1887a *Plataninium regulare* Felix, p. 148, pl. XXVII, f. 7.1889 *Plataninium regulare* Felix: Staub, p. 188.1890 *Plataninium regulare* Felix: Kaiser, p. 20.

Remarks: Originals in Mus. Ungar. Geol. Reichsanst.

Occurrence: Miocene (Lower Mediterranean); Hungary (Budafok).

Plataninium salinarum (Unger) Vater.1849 *Fegonium salinarum* Unger, p. 320, pl. XXXV, f. 25—27.1850 *Fegonium salinarum* Unger: Unger, p. 407.1872 *Fegonium salinarum* Unger: Schimper, p. 608.1873 *Fegonium salinarum* Unger: Stur, p. 9.1884 *Plataninium salinarum* (Unger) Vater, p. 837.1890 *Plataninium salinarum* (Unger): Kaiser, p. 20.

Occurrence: Tertiary; Galicia (Wieliczka).

Plataninium subaffine Vater.1884 *Plataninium subaffine* Vater, p. 843, pl. XXIX, f. 19—21.1890 *Plataninium subaffine* Vater: Kaiser, p. 20.

Occurrence: Derived from Lower Senonian; Brunswick (Helmstedt).

Plataninium vasculosum (Unger) Vater.1842 *Phegonium vasculosum* Unger, p. 101.1842a *Phegonium vasculosum* Unger: Unger, p. 173.1842b *Phegonium vasculosum* Unger: Unger, p. 748.

- 1845 *Fegonium vasculosum* Unger: Unger, p. 219.
 1847 *Fegonium vasculosum* Unger: Unger, p. 103, pl. XXVII, f. 7—9.
 1849 *Fegonium vasculosum* Unger: Unger, p. 321.
 1850 *Fegonium vasculosum* Unger: Unger, p. 407.
 1872 *Fegonium vasculosum* Unger: Schimper, p. 608.
 1884 *Plataninium vasculosum* (Unger) Vater, p. 837.
 1887a *Plataninium vasculosum* (Unger): Felix, p. 147.
 1890 *Plataninium vasculosum* (Unger): Kaiser, p. 20.
 Occurrence: Tertiary; Upper and Lower Austria, Styria, Galicia.

Plataninium sp.

- 1880 *Platanus aceroides* Goeppert: Schroeter, p. 35, pl. I, f. 10.
 1886 *Plataninium aceroides* (Goeppert) Windisch, p. 20.
 Remarks: Schroeter gave this wood a name instituted for leaves, but there is no evidence of connexion. The combination used by Windisch is also unacceptable, and as the wood is poorly preserved no specific name need be used. Original slides in the Geol. Dept., Brit. Mus. (Nat. Hist.).
 Occurrence: Tertiary; Arctic Canada (Mackenzie River).

? **Plataninium sp.**

- 1888 *Platanus* (?): Dawson, p. 33.
 Remarks: No description or figure. Requires confirmation.
 Occurrence: Upper Cretaceous (Belly River Series); Western Canada (Ribstone Creek).

Plataninium sp.

- 1894 *Plataninium sp.*: Felix, p. 102.
 Occurrence: Eocene (Sumgait Series); Caucasus (Apscheron).

Plataninium sp.

- 1930a *Plataninium*: Schönfeld, p. 70.
 Occurrence: ? Oligocene; Saxony (Deutzen).

Populocaulis Stopes & Fujii, 1910, p. 63.

Populocaulis yezoensis Stopes & Fujii.

- 1910 *Populocaulis yezoensis* Stopes & Fujii, p. 63, pl. VIII, f. 49.
 Remarks: Small twigs, transverse sections only. Probably unidentifiable. Originals in Geol. Dept., Brit. Mus. (Nat. Hist.).
 Occurrence: Upper Cretaceous; Japan (Hokkaido).

[**Populoxylon** Knowlton, 1899, p. 770.]

Remarks: Name mentioned, but rejected in favour of *Rhamnacinium* for a wood (*R. radiatum*) from the Yellowstone National Park. In any case the name would have to be regarded as a synonym of *Salicinium*, as most authorities agree that the woods of *Salix* and *Populus* cannot be separated.

Populus L.

See also *Salicinium* sp. and *Salix* sp.

? **Populus** sp.

1920 ? **Populus** sp.: Kräusel, p. 453, text-fig. 1—3.
Occurrence: Miocene; Silesia (Knurow).

[**Populus** sp. Dawson, 1875, p. 331.]

See **Rhamnacinium porcupinianum** and **R. triseriatum**.

Pritchardia Unger, 1842, p. 102.

[Non **Pritchardia** Seem. & Wendl., 1862, (Palmae).]

Pritchardia insignis Unger.

1842 **Pritchardia insignis** Unger, p. 102.
1842a **Pritchardia insignis** Unger: Unger, p. 177.
1845 **Pritchardia insignis** Unger: Unger, p. 260.
1850 **Pritchardia insignis** Unger: Unger, p. 522.
1890 **Pritchardia insignis** Unger: Kaiser, p. 36.
Occurrence: Tertiary; West Indies.

Pruninium Platen, 1908, p. 122.

Pruninium gummosum Platen.

1908 **Pruninium gummosum** Platen, p. 122, pl. III, f. 2—6.
Remarks: Compared with several living species of *Prunus*.
Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

Prunus L.**Prunus** sp.

1914 **Prunus** sp.: Szafer, p. 349.
Occurrence: Pleistocene; Poland.

Pterocarpus L.**Pterocarpus santalinus** L.

1890 **Pterocarpus santalinus** L.: Schenk, p. 902, f. 433.
Remarks: This wood is stated to be indistinguishable from that of the living red sandal-wood.
Occurrence: Age unknown; East Indies.

Quercinium Unger, 1842, p. 101.

Including **Kloedenia**, **Schimperites** and **Schmidites**.

Quercinium abromeiti Platen.

1908 **Quercinium abromeiti** Platen, p. 23.
Occurrence: Pliocene; California (Calistoga).

Quercinium anomalum Platen.

1908 **Quercinium anomalum** Platen, p. 47.
Occurrence: Miocene; California (Nevada County).

Quercinium astianum Pampaloni.

- 1904 *Quercinium astianum* Pampaloni, p. 540, f. 1—3.
 Occurrence: Pliocene; Piedmont (Ceresole d'Alba).

Quercinium austriacum Unger.

- 1842 *Quercinium austriacum* Unger, p. 101.
 1842a *Quercinium austriacum* Unger: Unger, p. 174.
 1842b *Quercinium austriacum* Unger: Unger, p. 748.
 1845 *Quercinium austriacum* Unger: Unger, p. 218.
 1847 *Quercinium austriacum* Unger: Unger, p. 107, pl. XXIX, f. 4—6.
 1850 *Quercinium austriacum* Unger: Unger, p. 405.
 1872 *Quercinium austriacum* Unger: Schimper, p. 662.
 1890 *Quercinium austriacum* Unger: Kaiser, p. 11.
 Occurrence: Tertiary; Austria (Bachmannig).

Quercinium böckhianum Felix.

- 1884 *Quercinium böckhianum* Felix, p. 21, pl. I, f. 6.
 1889 *Quercinium böckhianum* Felix: Staub, p. 187.
 1890 *Quercinium böckhianum* Felix: Kaiser, p. 11.
 Remarks: Very close to *Q. compactum*. Originals in Mus. Ungar. Geol. Reichsanst.
 Occurrence: Tertiary; Hungary (Medgyaszó).

Quercinium compactum Schleiden.

- 1855 *Quercinium compactum* Schleiden, p. 42.
 1883 *Quercinium compactum* Schleiden: Felix, p. 75, pl. II, f. 7.
 1884 *Quercinium compactum* Schleiden: Felix, p. 19.
 1889 *Quercinium compactum* Schleiden: Staub, p. 187.
 1890 *Quercinium compactum* Schleiden: Kaiser, p. 11.
 Remarks: Felix studied and figured Schleiden's original sections from the Jena Museum, and compared the species with *Quercus lusitana*.
 Occurrence: Tertiary; Hungary (Libethen).

Quercinium eocenicum Fritel & Viguiet.

- 1911 *Quercinium eocenicum* Fritel & Viguiet, p. 70, pl. I.
 Remarks: The authors note that this wood, although much older than that described by Eames (1910) shows no trace of the supposed primitive structure of the rays.
 Occurrence: Eocene (Sparnacian); Paris basin (Clairizet).

Quercinium helictoxyloides Felix.

- 1884 *Quercinium helictoxyloides* Felix, p. 17, pl. I, f. 3—5; IV, f. 3.
 1889 *Quercinium helictoxyloides* Felix: Staub, p. 187.
 1890 *Quercinium helictoxyloides* Felix: Kaiser, p. 12.
 Remarks: Root-wood. Originals in Mus. Ungar. Geol. Reichsanst.
 Occurrence: ? Pliocene; Hungary (Gyepüfűzes).

Quercinium knowltoni Felix.

- 1896 *Quercinium knowltoni* Felix, p. 250, pl. VI, f. 2.
 1899 *Quercinium knowltoni* Felix: Knowlton, p. 773.
 1910 *Quercinium knowltoni* Felix: Eames, p. 162.
 Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

Quercinium lamarensis Knowlton.

1899 *Quercinium lamarensis* Knowlton, p. 771, pl. CXVIII, f. 5; CXX, f. 2; CXXI, f. 1, 2.

1910 *Quercinium lamarensis* Knowlton: Eames, p. 162.

Remarks: Knowlton suggests that this may be identical with *Q. knowltoni* Felix.

Occurrence: Miocene; Yellowstone National Park.

Quercinium leptotichum (Schleiden) Felix.

1855 *Schimperites leptotichus* Schleiden, p. 42.

1883 *Quercinium leptotichum* (Schleiden) Felix, p. 77.

1884 *Quercinium leptotichum* (Schleiden): Felix, p. 23.

1889 *Quercinium leptotichum* (Schleiden): Staub, p. 188.

1890 *Quercinium leptotichum* (Schleiden): Kaiser, p. 12.

Remarks: Schleiden's original specimen was re-examined by Felix, and compared with various species of *Quercus*. It perhaps shows root structure.

Occurrence: Tertiary; Hungary (Libethen).

Quercinium lesquereuxi Platen.

1908 *Quercinium lesquereuxi* Platen, pp. 24, 50.

Occurrence: Miocene and Pliocene; California (Calistoga and Lake Tahoe).

Quercinium longiradiatum (Felix).

1887a *Quercinium staudi* var. *longiradiatum* Felix, p. 151, pl. XXVII, f. 3.

1890 *Quercinium staudi* var. *longiradiatum* Felix: Kaiser, p. 13.

Remarks: It is possible that this wood should be included in *Q. staudi*, but in any case the use of varietal names should be avoided. Probably root-wood. Original in Mus. Ungar. Geol. Reichsanst.

Occurrence: ? Tertiary; Hungary.

Quercinium montanum (Mercklin) Felix.

1855 *Quercinium rossicum* α *montanum* Mercklin, p. 28, pl. VII, f. 1—8.

1880 *Quercinium rossicum montanum* Mercklin: Krendowsky, p. 218 [Not seen].

1883 *Quercinium montanum* (Mercklin) Felix, p. 72, pl. III, f. 2, 7.

1890 *Quercinium montanum* (Mercklin): Kaiser, p. 12.

Remarks: Compared by Felix with the living *Q. tozza*, and by Mercklin with *Q. sessiliflora*.

Occurrence: Tertiary; Russia (Ekaterinoslav).

Quercinium pliocaenicum Schuster.

1908a *Quercinium pliocaenicum* Schuster, p. 52, pls. III, IV.

Remarks: Placed in the sub-genus *Erythrobalanus*. Schuster discusses the differences between his species and certain other fossil oaks: *Q. knowltoni*, *Q. böckhianum*, *Q. compactum*, and *Q. montanum*.

Occurrence: Pliocene; Idaho (Clover Creek).

[Quercinium primaevum (Goepfert).]

Synonym of *Quercinium quercoides*, q. v.

Quercinium quercoides (Goepfert).

- 1839 *Kloedenia quercoides* Goepfert, p. 521, pl. VIII B.
 1845 *Quercites primaevus* Goepfert, pp. 82, 84.
 1862 *Quercus primaeva* (Goepfert) Goepfert, p. 552.
 1876 *Quercites primaevus* Goepfert: Conwentz, p. 29.
 1881 *Quercus primaeva* (Goepfert): Goepfert, p. 133, pl. II, f. 6; III, f. 7.
 1883 *Quercites primaevus* Goepfert: Hofmann, p. 86.
 1883 *Quercinium primaevum* (Goepfert) Felix, p. 69, pl. III, f. 4, 8.
 1884 *Quercinium primaevum* (Goepfert): Felix, p. 12.
 1889 *Quercinium primaevum* (Goepfert): Staub, p. 187.
 1889 *Quercus primaeva* (Goepfert): Caspary, pp. 74—76.
 1890 *Quercites primaevus* Goepfert: Kaiser, p. 12.

Remarks: Since *Quercinium* is merely a form-genus for fossil wood closely resembling the recent *Quercus*, it would be pointless to revive the slightly earlier name *Kloedenia*, which Goepfert himself abandoned in 1845. There seems no reason, however, for adopting, as Felix did, the later specific name *Q. primaevum*, applied by Goepfert to a wood which may or may not be identical with *Q. quercoides*.

Felix 1883 (pp. 69—72) was able to show that one of Goepfert's originals came from Hungary (Tapolcsan), part of the same specimen being in Dresden and part in Berlin. The species is therefore really founded on this specimen as described by Felix, and records from other localities should be regarded as dubious. Felix compares the fossil with the recent *Quercus castaneaefolia*. Caspary (1889) also discusses the confusion which has centred around *Q. quercoides*, and compares his *Q. subgarryanum* with the wood described by Felix. Goepfert introduced the name *Quercites primaevus* for some wood from the amber-beds of the Baltic (stating that it was identical with his previously described *Kloedenia quercoides*) but these originals of Goepfert's seem to have been lost. Caspary re-examined one supposed original (he does not state the locality) and found that it showed no structure whatever. There is therefore no evidence concerning Goepfert's *Q. primaevus* from Samland, and the only described oak wood from the amber-beds is *Q. subgarryanum* Caspary.

Unger (1847, p. LXXIX, 1850, p. 405) cites *Quercites primaevus* as a synonym of his *Quercinium sabulosum*, and Conwentz cites the latter as a synonym of the former. *Q. sabulosum*, however, has never been adequately described, nor figured.

It is possible that *Q. vasculosum* may belong to this species.

Occurrence: Tertiary; Hungary (Tapolcsan). Also recorded from Samland, Silesia, Posen, Mecklenburg.

Quercinium rossicum Mercklin.

- 1855 *Quercinium rossicum* Mercklin, p. 27, pl. VI, f. 1—8.
 1865 *Quercinium rossicum* Mercklin: Eichwald, p. 57.
 1890 *Quercinium rossicum* Mercklin: Kaiser, p. 13.

Remarks: Compared with *Quercus pedunculata*. For *Q. rossicum* a *montanum*, see *Q. montanum*.

Occurrence: Tertiary; Russia (Alexandrovsk in Ekaterinoslav).

Quercinium sabulosum Unger.

- 1842 *Quercinium sabulosum* Unger, p. 101.
 1842a *Quercinium sabulosum* Unger: Unger, p. 173.
 1842b *Quercinium sabulosum* Unger: Unger, p. 747.
 1845 *Quercinium sabulosum* Unger: Unger, p. 218.
 1847 *Quercinium sabulosum* Unger: Unger, p. 108.

1850 *Quercinium sabulosum* Unger: Unger, p. 405.

1872 *Quercinium sabulosum* Unger: Schimper, p. 661.

1889 *Quercinium sabulosum* Unger: Staub, p. 188.

1890 *Quercinium sabulosum* Unger: Kaiser, p. 13.

Remarks: Unger (1845) cites Goeppert's *Kloedenia quercoides* and *Quercites primaevus* as synonyms of *Q. sabulosum*, without giving any reasons. Conwentz (1876) includes *Q. sabulosum* as a synonym of *Quercites primaevus*.

The species never seems to have been figured, and the description is inadequate.

Occurrence: Tertiary; various localities in Austria, Moravia, Hungary, Silesia.

Quercinium solerederi Platen.

1908 *Quercinium solerederi* Platen, p. 41.

Occurrence: Mio-Pliocene; California.

Quercinium staubi Felix.

1884 *Quercinium staubi* Felix, p. 15, pl. I, f. 2.

1889 *Quercinium staubi* Felix: Staub, p. 187.

1890 *Quercinium staubi* Felix: Kaiser, p. 13.

Remarks: Resembles *Q. quercoides* and the recent *Quercus castaneaeifolia*. Originals in Mus. Ungar. Geol. Reichsanst. For *Q. staubi* var. *longiradiatum*, see *Q. longiradiatum*.

Occurrence: Pliocene; Hungary (Gyepüfüzes).

Quercinium subgarryanum (Caspary).

1888 *Quercus subgarryana* Caspary, p. 44.

1889 *Quercus subgarryana* Caspary: Caspary, p. 71, pl. XIII, f. 6—11.

1890 *Quercus subgarryana* Caspary: Kaiser, p. 14.

Remarks: Compared with the North American *Quercus garryana*. Originals in Mus. Bot. Gart., Königsberg. Caspary points out the differences between this wood and the *Quercinium* "*primaevum*" described by Felix; Goeppert had recorded *Quercites primaevus* (as a synonym of his *Kloedenia quercoides*) from the amber-beds of the Baltic, but there does not appear to be any evidence as to what this amber-bed wood really was. Felix's re-description of Goeppert's species was partly founded on some originals of Goeppert's from Hungary. (See also the remarks on *Quercinium quercoides*).

Occurrence: Tertiary; Königsberg.

Quercinium transiens (Conwentz).

1876 *Quercites transiens* Conwentz, p. 30.

1890 *Quercites transiens* Conwentz: Kaiser, p. 14.

Occurrence: Erratic block; Galicia.

Quercinium transylvanicum Unger.

1842 *Quercinium transylvanicum* Unger, p. 101.

1842a *Quercinium transylvanicum* Unger: Unger, p. 174.

1845 *Quercinium transylvanicum* Unger, p. 218.

1847 *Quercinium transylvanicum* Unger, p. 103.

1850 *Quercinium transylvanicum* Unger, p. 405.

1872 *Quercinium transylvanicum* Unger: Schimper, p. 662.

1889 *Quercinium transylvanicum* Unger: Staub, p. 188.

1890 *Quercinium transylvanicum* Unger: Kaiser, p. 14.

Occurrence: Tertiary; Transylvania.

Quercinium vasculosum (Schleiden) Felix.1855 *Schmidites vasculosus* Schleiden, p. 39.1883 *Quercinium vasculosum* (Schleiden) Felix, p. 76, pl. II, f. 2.1884 *Quercinium vasculosum* (Schleiden): Felix, p. 20.1889 *Quercinium vasculosum* (Schleiden): Staub, p. 187.1890 *Quercinium vasculosum* (Schleiden): Kaiser, p. 14.

Remarks: Felix examined the originals in the Jena Museum and showed that the wood belonged to *Quercinium*, and not to the *Leguminosae*, as Schleiden thought. Felix puts it near *Q. quercoides*.

Occurrence: Tertiary; Hungary (Tapolcsan).

Quercinium wardi Platen.1908 *Quercinium wardi* Platen, p. 49.

Occurrence: Mio-Pliocene; California.

Quercinium sp.

1910 "Fossil oak": Eames, p. 161, pl. VIII, f. 5, 6.

Remarks: The author discusses the structure of the broad rays of an unnamed Miocene oak, and its bearing on the origin of the broad rays in *Quercus*. See also remarks on *Q. eocenicum*.

Occurrence: Miocene; California (gold gravels).

Quercinium sp.1912 *Quercinium* sp.: Becke, p. 85.

Occurrence: Tertiary; Germany (Joachimstal).

Quercinium sp.

1885 Eichenholz: Quenstedt, p. 1150, pl. XCVIII, f. 23.

Remarks: Two poor figures of a silicified oak wood; horizon and locality not stated.

[Quercites primaevus Goeppert.]

Synonym of *Quercinium quercoides*, q. v.

[Quercoxylon Hofmann, 1929, p. 82.]

Synonym of *Quercinium*. See *Quercus cerris*.

Quercus L.**Quercus cf. cerris** L.1929 *Quercoxylon cerris* Hofmann, p. 82.

Remarks: No figure nor description. See remarks on *Fraxinus excelsior*.

Occurrence: Plio-Pleistocene; Hungary (Csadberge).

[Quercus primaeva Goeppert.]

Synonym of *Quercinium quercoides*, q. v.

Quercus sp.1873 *Quercus* sp.: Dawson, p. 68.

Remarks: Requires confirmation.

Occurrence: Cretaceous; British Columbia.

Quercus sp.1914 *Quercus* sp.: Szafer, p. 349.1926a *Quercus*: Fietz, p. 417.

Occurrence: Pleistocene; Poland.

Prehistoric (Loess); Czechoslovakia (Brno).

Reynosia Grisebach.**Reynosia texana** Penhallow.1907 *Reynosia texana* Penhallow, p. 97, f. 4, 5.1924 *Reynosia texana* Penhallow: Berry, p. 75.Remarks: Said to resemble the existing *R. latifolia* of Florida and Cuba. Originals in Peter Redpath Mus., Montreal.

Occurrence: Eocene (Yegua); Texas (Somerville).

Reveesia Walp.**Reveesia wallichii** R. Br.1911a *Reveesia wallichii* R. Br.: Schuster, p. 246.

Remarks: The identification should be accepted with reserve.

Occurrence: Pithecanthropus Beds; Java.

Rhamnacinium Felix, 1894, p. 89.**Rhamnacinium affine** Felix.1894 *Rhamnacinium affine* Felix, p. 88, pl. VIII, f. 3a—d.Remarks: Compared with the living *Prinos* and *Pomaderris*, and the fossil *Rhamnacinium primaevum*.

Occurrence: ? Eocene (Sumgait series); Caucasus (Apscheron).

Rhamnacinium porcupinianum Penhallow.1875 *Populus* sp. (pars): Dawson, p. 331.1903 *Rhamnacinium porcupinianum* Penhallow, p. 48, f. 14—16, 21, 22.

Occurrence: Eocene; Saskatchewan (Porcupine Creek).

Rhamnacinium primaevum (Caspary) Felix.1888 *Schinus primaevum* Caspary, p. 39.1889 *Schinus primaevum* Caspary: Caspary, p. 20, pl. IV, f. 13; V, f. 1—4.1890 *Schinus primaevum* Caspary: Kaiser, p. 26.1894 *Rhamnacinium primaevum* (Caspary) Felix, pp. 87, 89.Remarks: Compared by Caspary with *Schinus molle* (*Anacardiaceae*), but referred by Felix to the *Rhamnaceae* and compared with *Prinos* and *Pomaderris*.

Occurrence: ? Tertiary; West Prussia (Pempau).

Rhamnacinium radiatum Felix.1896 **Rhamnacinium radiatum** Felix, p. 252, pl. VI, f. 3.1899 **Rhamnacinium radiatum** Knowlton, p. 769, pl. CXVIII, f. 6, 7; CXIX, f. 1.Remarks: Felix notes the distinctions from *R. affine* and *R. primaevum*.

Occurrence: Miocene; Yellowstone National Park (Amethyst Mt.).

Rhamnacinium texanum Penhallow.1907 **Rhamnacinium texanum** Penhallow, p. 96, f. 1—3.1924 **Rhamnacinium texanum** Penhallow; Berry, p. 74.

Remarks: Originals in Peter Redpath Museum, Montreal.

Occurrence: Eocene (Yegua); Texas (Somerville).

Rhamnacinium triseriatum Penhallow.1875 **Populus** sp. (pars): Dawson, p. 331.1903 **Rhamnacinium triseriatum** Penhallow, p. 54, f. 17—20.

Occurrence: Eocene; Saskatchewan (Porcupine Creek).

Rhamnus L.**Rhamnus cathartica** L.1926a **Rhamnus cathartica** L.: Fietz, p. 419.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

[Rhizoalnoxylon Conwentz, 1880.]Included in **Alnoxylon** Felix.**Rhodium** Unger, 1850, p. 475.**Rhodium juglandinum** Unger.1850 **Rhodium juglandinum** Unger, p. 475.1889 **Rhodium juglandinum** Unger: Staub, p. 190.1890 **Rhodium juglandinum** Unger: Kaiser, p. 26.

Occurrence: Tertiary; Hungary (Antal near Schemnitz).

Rhodium philippinense Crié.1889 **Rhodium philippinense** Crié, p. 86, pl. IX (XVII), f. 3, 4.

Occurrence: Tertiary; Philippines (Luzon, Manilla, S. Juan del Monte).

Rhodium ungeri Mercklin.1855 **Rhodium ungeri** Mercklin, p. 21, pl. I, f. 1, 2; II, f. 1—10.1865 **Rhodium ungeri** Mercklin: Eichwald, p. 65.1890 **Rhodium ungeri** Mercklin: Kaiser, p. 26.Remarks: Compared with *Rhus juglandifolia*.

Occurrence: ? Cretaceous; Russia (Durasovka in Saratov).

[Roemeria Unger, 1852a, p. 95.]Remarks: Non *Roemeria* Medicus, 1792. For *R. americana* Unger see *Dryoxylon americanum*.

Rohlfisia Schenk, 1880, p. 660.**Rohlfisia celastroides** Schenk.

- 1880 **Rohlfisia celastroides** Schenk, p. 660.
 1883 **Rohlfisia celastroides** Schenk: Schenk, p. 9, pl. IV, f. 12.
 1888 **Rohlfisia celastroides** Schenk: Schenk, p. 20.
 1890 **Rohlfisia celastroides** Schenk: Kaiser, p. 27.

Remarks: Probably *Celastraceae*.

Occurrence: Upper Cretaceous; Libyan Desert.

Rosthornia Unger, 1842, p. 101.**Rosthornia carinthiaca** Unger.

- 1842 **Rosthornia carinthiaca** Unger, p. 101.
 1842a **Rosthornia carinthiaca** Unger: Unger, p. 175.
 1845 **Rosthornia carinthiaca** Unger: Unger, p. 225.
 1850 **Rosthornia carinthiaca** Unger: Unger, p. 421.
 1890 **Rosthornia carinthiaca** Unger: Kaiser, p. 16.

Remarks: Referred doubtfully to the *Salicaceae* by Unger. Lignier (1907, p. 271 footnote) suggests that it might possibly belong to the *Magnoliaceae*.

Occurrence: Upper Cretaceous (Gosau formation); Carinthia.

Ruyschioxylon Hofmann, 1884, p. 183.**Ruyschioxylon sumatrense** Hofmann.

- 1884 **Ruyschioxylon sumatrense** Hofmann, p. 183 (32).
 1890 **Ruyschioxylon sumatrense** Hofmann: Kaiser, p. 24.
 1930 **Ruyschioxylon sumatrense** Hofmann: Schönfeld, p. 124.

Remarks: A silicified liane referred to the *Ternstroemiaceae*, and compared with the recent *Ruyschia*. According to Schönfeld it is very close to *Helictoxylon schenki*. Original in Coll. Univ. Utrecht.

Occurrence: Tertiary; Sumatra.

Sabiocaulis Stopes & Fujii, 1910, p. 66.**Sabiocaulis sakurii** Stopes & Fujii.

- 1910 **Sabiocaulis sakurii** Stopes & Fujii, p. 66, pl. VIII, f. 54; IX, f. 55—57.

Remarks: Compared with *Sabia japonica*. Some of the originals are in the Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Upper Cretaceous; Japan (Hokkaido).

Sabulia Stopes, 1912, p. 93.**Sabulia scottii** Stopes.

- 1912 **Sabulia scottii** Stopes, p. 93, pl. VI, f. 2; VIII, f. 9.
 1915 **Sabulia scottii** Stopes: Stopes, p. 272, text-fig. 82—84.
 Remarks: Affinities unknown. Originals in Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Lower Cretaceous (Aptian); England (Woburn).

Salicinium Unger, 1850, p. 420.**Salicinium antiquum** (Falqui).1906 *Salix purpurea* Linn. β *antiqua* Falqui, p. 21, pl. I, f. 6.

Remarks: Presuming that this belongs to the *Salicaceae*, it should obviously be put in the form-genus *Salicinium*, and not identified with a living species.

Occurrence: Miocene; Sardinia.

Salicinium biradiatum (Lignier).1907 *Salicinoxylon biradiatum* Lignier, p. 272, pl. XVIII, f. 18—24; XXIII, f. 84—92.

Remarks: This wood is referred with some hesitation to the *Salicaceae*.

Occurrence: Cenomanian; Seine-Inférieure (La Hève).

Salicinium bruxellense Hofmann.1884 *Salicinium bruxellense* Hofmann, p. 193 (42).1890 *Salicinium bruxellense* Hofmann: Kaiser, p. 16.

Remarks: Original in Coll. Univ. Utrecht.

Occurrence: Lower Tertiary; Belgium (near Brussels).

Salicinium messinianum Pampaloni.1904 *Salicinium messinianum* Pampaloni, p. 545, f. 10—11.

Occurrence: Pliocene; Piedmont.

Salicinium miocenicum (Kaiser).1880a *Salicinoxylon miocenicum* Kaiser, p. 511.1890 *Salicinoxylon miocenicum* Kaiser: Kaiser, p. 16.

Remarks: Kaiser apparently overlooked *Salicinium populinum* Unger when he stated that no fossil salicineous wood had previously been described. Kaiser's name *Salicinoxylon* is superfluous.

See *Salix* ? sp. for a record by Fietz from the Pleistocene of Silesia.

Occurrence: Probably Miocene; Schleswig-Holstein (Sylt).

Salicinium populinum Unger.1850 *Salicinium populinum* Unger, p. 420.1882 *Salicinium populinum* Unger: Felix, p. 36.1890 *Salicinium populinum* Unger: Kaiser, p. 16.

Occurrence: ? Tertiary; Vienna.

Salicinium varians Hofmann.1884 *Salicinium varians* Hofmann, p. 191.1890 *Salicinium varians* Hofmann: Kaiser, p. 16.

Remarks: Caspary (1889, p. 5) doubts whether this wood belongs to the *Salicaceae*. Originals in Coll. Univ. Utrecht.

Occurrence: Senonian; Holland (Pietersberg near Maastricht).

? **Salicinium** sp.1873 *Populus* sp.: Dawson, p. 68.1888 *Populus*: Dawson, p. 33.

Remarks: No description or illustration. Other woods referred by Dawson to *Populus* were described by Penhallow as *Rhamnacinium*.

Occurrence: Upper Cretaceous; British Columbia (Vancouver I.), Western Canada (Swift Current).

[**Salicinoxyton** Kaiser, 1880a, p. 511.]

Synonym of **Salicinium**, q. v. for all species described under **Salicinoxyton**.

Salix L.

[**Salix purpurea** L. β *antiqua* Falqui.]

See **Salicinium antiquum**.

Salix sp.

1914 **Salix** sp.: Szafer, p. 347.

Occurrence: Pleistocene; Poland.

Salix ? sp.

1926 **Salicinoxyton miocenicum** Kaiser: Fietz, p. 234.

1926a **Salix** ? : Fietz, p. 417.

Remarks: There does not seem sufficient reason for referring a Pleistocene wood to Kaiser's species, unless indeed *S. miocenicum* is to cover every fossil wood belonging to the *Salicaceae*. The name *Salix* is queried because, as Fietz points out, there is no absolute distinction between the woods of *Salix* and of *Populus*. (See remarks on nomenclature in the introduction).

Occurrence: Pleistocene; Silesia (Weidenau), Czechoslovakia (Brno).

Santalum L.

Santalum sp.

1882b **Santalum** sp.: Kraus, p. 8.

Remarks: The supposed former occurrence of *S. album* on Juan Fernandez is so far confirmed that Kraus regards some wood investigated by him as undoubtedly belonging to the genus *Santalum*, which is today confined to the East Indies.

Occurrence: Sub-fossil; Juan Fernandez.

[**Sapindopsoxyton** Pfeiffer & van Heurn, 1928, p. 1009.]

Included in **Sapindoxylon**, q. v. for **S. klitzingi**.

Sapindoxylon Kräusel, 1922, p. 256.

Sapindoxylon janssonii Kräusel.

1922 **Sapindoxylon janssonii** Kräusel, p. 256, pl. I, f. 9; II, f. 3; III, f. 6; V, f. 5; VI, f. 2; text-fig. 22.

1922 ? **Sapindoxylon janssonii** Kräusel: Kräusel, p. 258, pl. I, f. 8.

1928 **Sapindoxylon janssonii** Kräusel: Pfeiffer & van Heurn, p. 1007.

Remarks: Compared particularly with the genus *Pometia*.

Occurrence: Tertiary; Java, South Sumatra.

Sapindoxylon klitzingi (Pfeiffer & van Heurn).

1928 **Sapindopsoxylon klitzingi** Pfeiffer & van Heurn, p. 1005, pl. I, f. 1.

Remarks: Referred to the *Sapindaceae*, but considered to differ from Kräusel's *Sapindoxylon* in certain respects. The authors disclaim any intention of expressing a relationship with *Sapindopsis* Fontaine. The new generic name seems to me both unfortunate and unnecessary.

Occurrence: Tertiary; Java (Bolang).

Sapotoxylon Felix, 1882.

Sapotoxylon gümbelii Felix.

1882 **Sapotoxylon gümbelii** Felix, p. 54.

1883 **Sapotoxylon gümbelii** Felix: Felix, p. 67, pl. II, f. 5, 8.

1890 **Sapotoxylon gümbelii** Felix: Kaiser, p. 33.

Remarks: Silicified wood, apparently a water-worn block. Referred to the Sapotaceae, but Kaiser notes that it is also similar to the wood of the Anonaceae.

Occurrence: Tertiary?; Wagenhofen near Neuburg on the Danube.

Sapotoxylon taeniatum Felix.

1882 **Sapotoxylon taeniatum** Felix, p. 56.

1883 **Sapotoxylon taeniatum** Felix: Felix, p. 68, pl. III, f. 5, 6.

1890 **Sapotoxylon taeniatum** Felix: Kaiser, p. 33.

Occurrence: Locality and horizon unknown. Original in Munich Museum.

Sauruopsis Stopes & Fujii, 1910, p. 58.

Sauruopsis niponensis Stopes & Fujii.

1910 **Sauruopsis niponensis** Stopes & Fujii, p. 58, pl. VII, f. 42—47, text-fig. 19.

Remarks: A small stem believed to belong to the *Saururaceae*. Some of the originals are in the Geol. Dept., Brit. Mus. (Nat. Hist.).

Occurrence: Upper Cretaceous; Japan (Hokkaido).

[**Schimperites** Schleiden, 1855, p. 42.]

Synonym of **Quercinium**.

[**Schimperites leptotichus** Schleiden.]

Synonym of **Quercinium leptotichum**, q. v.

[**Schinus primaevum** Caspary.]

Synonym of **Rhamnacinium primaevum**, q. v.

Schleidenites Unger, 1842, p. 102.

Schleidenites compositus Unger.

1842 **Schleidenites compositus** Unger, p. 102.

1842a **Schleidenites compositus** Unger: Unger, p. 178.

1845 **Schleidenites compositus** Unger: Unger, p. 266.

1850 *Schleidenites compositus* Unger: Unger, p. 526.

1889 *Schleidenites compositus* Unger: Staub, p. 191.

1890 *Schleidenites compositus* Unger: Kaiser, p. 32.

Remarks: Referred by Unger to the *Leguminosae*.

Occurrence: Tertiary; Hungary.

[**Schmidites** Schleiden, 1855, p. 39.]

Synonym of *Quercinium*.

[**Schmidites vasculosus** Schleiden.]

Synonym of *Quercinium vasculosum*, q. v.

Schmiedeliopsis Felix, 1882, p. 72.

Schmiedeliopsis zirkelii Felix.

1882 *Schmiedeliopsis zirkelii* Felix, p. 72.

1883a *Schmiedeliopsis zirkelii* Felix: Felix, p. 16, pl. II, f. 6, 8.

1890 *Schmiedeliopsis zirkelii* Felix: Kaiser, p. 26.

Remarks: Possibly belongs to the *Sapindaceae*, as the wood resembles that of *Schmiedelia haemorrhoea*. Original in Munich.

Occurrence: Tertiary; Antigua.

[**Shoreoxylon** den Berger, 1923, p. 147 & 1927, p. 497.]

Remarks: For Dipterocarpaceous woods resembling those of the *Shoreae* (*Dipterocarpoxyton* in part). Kräusel (1925, 1926) doubts whether it is yet possible satisfactorily to subdivide the genus *Dipterocarpoxyton*, and for the purposes of this catalogue I have followed Kräusel.

[**Shoreoxylon djambiense**, *S. kräuseli*, *S. moroides*.]

See *Dipterocarpoxyton*.

[**Shoreoxylon palembangense** (Kräusel) den Berger.]

See *Caesalpinioxylon*.

Sillimannia Unger, 1850, p. 524.

Sillimannia texana Unger.

1850 *Sillimannia texana* Unger, p. 524.

1852a *Sillimannia texana* Unger: Unger, p. 95.

1890 *Sillimannia texana* Unger: Kaiser, p. 36.

Remarks: Referred by Unger to the *Sterculiaceae*.

Occurrence: Cretaceous; Texas.

Simarubinium Platen, 1908, p. 54.

Simarubinium crystallophorum Platen.

1908 *Simarubinium crystallophorum* Platen, p. 54.

Occurrence: Tertiary; California.

Simarubinium engelhardti Platen.1908 **Simarubinium engelhardti** Platen, p. 56.

Occurrence: Tertiary; California.

Sjögrenia Felix, 1894, p. 99.**Sjögrenia crystallophora** Felix.1894 **Sjögrenia crystallophora** Felix, p. 93, pl. IX, f. 1a—d, 2a—b.Remarks: Possibly related to the *Aurantiaceae*.

Occurrence: Eocene (Sumgait Series); Caucasus (Baku).

Staphylea L.**Staphylea** sp.1926a **Staphylea**: Fietz, p. 418.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Staubia Felix, 1884, p. 28.**Staubia eriodendroides** Felix.1884 **Staubia eriodendroides** Felix, p. 29, pl. II, f. 2, 4—6, 8.1889 **Staubia eriodendroides** Felix: Staub, p. 190.1890 **Staubia eriodendroides** Felix: Kaiser, p. 25.Remarks: The genus *Staubia* is stated to be close to *Dombeyoxylon*, and *S. eriodendroides* to be intermediate between *Eriodendron* and *Pterospermum*. Originals in Min. Mus. Dresden.

Occurrence: Tertiary; Hungary.

Stephanoxylon Felix, 1882, p. 43.**Stephanoxylon dubium** Felix.1882 **Stephanoxylon dubium** Felix, p. 43.1890 **Stephanoxylon dubium** Felix: Kaiser, p. 36.

Remarks: Silicified wood of uncertain systematic position. Not figured. Original in Munich.

Occurrence: ? Tertiary; ? Cracow.

Suevioxylon Kräusel, 1928, p. 253.**Suevioxylon zonatum** Kräusel.1928 **Suevioxylon zonatum** Kräusel, p. 253, f. 5—8.

Remarks: The wood is unidentifiable and indeed is so poorly preserved that its reference to the Dicotyledones cannot be regarded as fully established.

Occurrence: Jurassic (Braunjura β); Germany (Heubach).**Sumatroxylon** den Berger, 1923, p. 145.**Sumatroxylon mollii** (Kräusel) den Berger.1922 **Anacardioxylon mollii** Kräusel, p. 252, pl. II, f. 5; IV, f. 4, 5; V, f. 2—4; VI, f. 3, 6, 7; VII, f. 3—5.

1923 *Sumatroxylon mollii* (Kräusel) den Berger, p. 145.

1925 *Sumatroxylon mollii* (Kräusel): Kräusel, p. 340.

Remarks: Kräusel at first referred this wood to the *Anacardiaceae*, but afterwards accepted the neutral name suggested by den Berger, who compared it with the *Burseraceae*. The name *Dryoxylon* might well have been used.

Occurrence: Miocene; Sumatra.

Swietenioxylon Hofmann, 1883, p. 105.

Swietenioxylon sternbergense Hofmann.

1883 *Swietenioxylon sternbergense* Hofmann, p. 105.

1890 *Swietenioxylon sternbergense* Hofmann: Kaiser, p. 25.

Remarks: *Meliaceae*. Said to resemble *Swietenia mahagoni*.

Occurrence: Upper Oligocene; Mecklenburg (Sternberg).

Taenioxylon Felix, 1882, p. 64.

Taenioxylon blanfordi Felix.

1883a *Taenioxylon blanfordi* Felix, p. 13, pl. I, f. 5, 6.

1890 *Taenioxylon blanfordi* Felix: Kaiser, p. 37.

Occurrence: ? Tertiary; "Ost-Indien".

Taenioxylon eperuoides Felix.

1886 *Taenioxylon eperuoides* Felix, p. 491, pl. XII, f. 5, 6.

1890 *Taenioxylon eperuoides* Felix: Kaiser, p. 37.

Remarks: Compared with the living *Eperua decandra* (*Caesalpiniaceae*). Originals in Min. Mus. Dresden.

Occurrence: ? Pliocene; Philippines (Valentia).

Taenioxylon ingaeforme Felix.

1882 *Taenioxylon ingaeforme* Felix, p. 79.

1883a *Taenioxylon ingaeforme* Felix: Felix, p. 12, pl. I, f. 7—9.

1890 *Taenioxylon ingaeforme* Felix: Kaiser, p. 37.

Remarks: Probably *Mimoseae*; resembles *Inga fastuosa*.

Occurrence: ? Tertiary; Brazil.

Taenioxylon irregulare Felix.

1882 *Taenioxylon irregulare* Felix, p. 65.

1883a *Taenioxylon irregulare* Felix: Felix, p. 11, pl. I, f. 1, 2.

1890 *Taenioxylon irregulare* Felix: Kaiser, p. 37.

Remarks: Systematic position uncertain.

Occurrence: Tertiary; Antigua.

Taenioxylon multiradiatum Felix.

1883a *Taenioxylon multiradiatum* Felix, p. 11, pl. I, f. 10, 11; II, f. 10.

1890 *Taenioxylon multiradiatum* Felix: Kaiser, p. 37.

1918 *Taenioxylon multiradiatum* Felix: Berry, p. 30, pls. XIV, XV.

Remarks: Probably *Papilionaceae*.

Occurrence: Tertiary; Antigua.

Oligocene; Canal Zone.

[*Taenioxylon pacificum* Crié.]1889 *Taenioxylon pacificum* Crié, p. 90.

Remarks: Nomen nudum.

Occurrence: Pliocene; Molucca (Halmahera).

Taenioxylon pannonicum Felix.1887a *Taenioxylon pannonicum* Felix, p. 145, pl. XXVII, f. 1, 2.1889 *Taenioxylon pannonicum* Felix: Staub, p. 190.1890 *Taenioxylon pannonicum* Felix: Kaiser, p. 37.

Remarks: Originals in Mus. Ungar. Geol. Reichsanst.

Occurrence: ? Tertiary; Hungary.

Taenioxylon porosum Felix.1894 *Taenioxylon porosum* Felix, p. 103, pl. X, f. 3.Remarks: Felix suggests that perhaps this wood ought to be put in *Sapotoxylon*.

Occurrence: Eocene (Sumgait Series); Caucasus (Apscheron).

Taenioxylon varians Felix.1882 *Taenioxylon varians* Felix, p. 64.1883a *Taenioxylon varians* Felix: Felix, p. 10, pl. I, f. 3, 4.1884 *Taenioxylon varians* Felix: Vater, p. 852.1890 *Taenioxylon varians* Felix: Kaiser, p. 37.

Remarks: Systematic position uncertain.

Occurrence: Lower Senonian; Germany (Harzburg).
Tertiary; Antigua.*Taenioxylon* sp.1884 *Taenioxylon* sp.: Vater, p. 852.1890 *Taenioxylon* sp.: Kaiser, p. 38.Remarks: Resembles *T. varians*.

Occurrence: Lower Senonian; Brunswick (Helmstedt).

Tarrietioxylon Kräusel, 1922, p. 259.*Tarrietioxylon sumatrense* Kräusel.1922 *Tarrietioxylon sumatrense* Kräusel, p. 259, pl. II, f. 4; IV, f. 2, 3, 6; VI, f. 4, 5, 9; text-fig. 23.Remarks: Referred to the *Sterculiaceae* and compared with *Tarrietia*.

Occurrence: Tertiary; South Sumatra.

Tchihatcheffites Unger, 1866a, p. 324.[= *Tchihatchewia* Unger, 1863, p. 516, preoccupied.]*Tchihatcheffites byzantina* Unger.1863 *Tchihatchewia byzantina* Unger in *Tchihatchef*, p. 516.1866a *Tchihatcheffites byzantina* (Unger) Unger, p. 324, pl. XVII, f. 3, 4.

Remarks: Family unknown (Unger).

Occurrence: Tertiary; Thrace (Lake Derkos).

Ternstroemiacinium Felix, 1894, p. 100.

Ternstroemiacinium euryoides Felix.

1894 **Ternstroemiacinium euryoides** Felix, p. 99, pl. X, f. 4.

1911 **Ternstroemiacinium euryoides** Felix: Tuzson, p. 49.

1930 **Ternstroemiacinium euryoides** Felix: Schönfeld, p. 124.

Remarks: Compared with *Eurya latifolia*.

Occurrence: Eocene (Sumgait Series); Caucasus (Apscheron).

Ternstroemiacinium kräuseli (Schönfeld).

1930 **Ternstroemioxylon kräuseli** Schönfeld, p. 119, f. 10—18.

Remarks: Schönfeld gives no reason for substituting a new name for the perfectly adequate form-genus *Ternstroemiacinium* Felix.

Occurrence: Miocene; Hessen (Lauterbach am Vogelsberg).

[**Ternstroemioxylon** Schönfeld, 1930.]

Synonym of **Ternstroemiacinium** Felix.

Tetranthera Jacq.

Tetranthera alnoides Miq.

1911a **Tetranthera alnoides** Miq.: Schuster, p. 246, f. 2, 3.

Remarks: The identification should be accepted with reserve.

Occurrence: Pithecanthropus Beds; Java.

Tilia L.

Tilia sp.

1926a **Tilia**: Fietz, p. 419.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Tilioxylon Hofmann, 1929, p. 82.

Tilioxylon sp.

1928 **Tilia** sp.: Hofmann, p. 3, pl. 1, f. 2.

1929 **Tilioxylon** sp.: Hofmann, p. 82.

Occurrence: Pliocene or Pleistocene; Hungary (Csadberge).

[**Trochodendromagnolia** Zander.]

1923 **Trochodendromagnolia**: Zander, p. 40, 41, pl. III.

Remarks: The above name refers to some isolated tracheids obtained by macerating an Eocene lignite. No species is mentioned, and the name was apparently used to suggest a somewhat hypothetical intermediate form. It is not clear whether all the tracheids figured are supposed to have come from the same plant, and it seems very unlikely that such material could ever be definitely identified.

Occurrence: Eocene; Saxony (Geiseltal).

Ulminium Unger, 1842, p. 101.**Ulminium columbianum** (Penhallow) Nagelhard.1907a *Ulmus columbiana* Penhallow, p. 299, pl. VIII.1908 *Ulmus* "columbiensis" Penhallow: Penhallow, p. 93.1922 *Ulminium columbianum* (Penhallow) Nagelhard, p. 19.

Remarks: Penhallow states that he first regarded this wood as *Rhamnacinium*, but that a more critical examination showed it to be an *Ulmus* not readily assignable to a known type. Under the circumstances a more cautious generic reference seems desirable.

Occurrence: Oligocene; British Columbia (Midway).

[Ulminium diluviale Unger.]

Synonym of *Laurinium diluviale*, q. v.

Ulminium hungaricum (Lingelsheim).1917 *Ulmoxylon hungaricum* Lingelsheim, p. 561, f. 12, 13.

Remarks: Type in Mus. Geol. Reichsanst. Budapest.

Occurrence: Miocene; Hungary (Beocsin).

Ulminium kleinii (Tuzson).1909 *Celtitis kleinii* Tuzson, p. 376. (Nomen).1911 *Celtites kleinii* Tuzson, p. 50, pl. I, f. 3; text-fig. 22—25.

Occurrence: Tertiary (Schotter Beds); Hungary (Lake Balaton).

Ulminium lapidariorum (Unger).1842 *Cottaites lapidariorum* Unger, p. 102.1842a *Cottaites lapidariorum* Unger: Unger, p. 176.1845 *Cottaites lapidariorum* Unger: Unger, p. 265.1850 *Cottaites lapidariorum* Unger: Unger, p. 526.1854 *Cottaites lapidariorum* Unger: Unger, p. 182, pl. VII, f. 1—3.1879 *Ulmoxylon* sp.: Kaiser, p. 100.1883 *Ulmoxylon lapidariorum* (Unger) Felix, p. 63.1890 *Ulmoxylon lapidariorum* (Unger): Kaiser, p. 18.1908 *Ulmoxylon lapidariorum* (Unger): Platen, p. 27.

Remarks: Kaiser (1879) described a silicified wood from Gleichenberg as *Ulmoxylon*, without giving a specific name. He showed that Unger's *Cottaites lapidariorum* belonged to the *Ulmaceae* and not the *Leguminoseae*, and stated that it was very probably synonymous with his *Ulmoxylon*.

Occurrence: Miocene; Styria (Gleichenberg).

Ulminium lovisatoi (Falqui).1906 *Ulmoxylon lovisatoi* Falqui, p. 16, pl. I, f. 4.

Remarks: Lingelsheim (1917, p. 563) doubts whether this wood has anything to do with *Ulmus*. See addendum, p. 90.

Occurrence: Miocene; Sardinia.

Ulminium pliocenicum Pampaloni.1904 *Ulminium pliocenicum* Pampaloni, p. 542, f. 4—6.

Occurrence: Pliocene; Piedmont (Ceresole d'Alba).

Ulmium protoamericanum (Penhallow) Nagelhard.

- 1907a *Ulmus protoamericana* Penhallow, p. 298, pl. VII.
 1908 *Ulmus protoamericana* Penhallow: Penhallow, p. 94.
 1922 *Ulmium protoamericanum* (Penhallow) Nagelhard, p. 20.
 Remarks: Penhallow (1907b, p. 449) states that a wood close to
U. americana occurs in the Pleistocene of Elmira, N. Y.
 Occurrence: Oligocene; British Columbia (Kettle River).

Ulmium protoracemosum (Penhallow) Nagelhard.

- 1907a *Ulmus protoracemosa* Penhallow, p. 297, pls. IV—VI.
 1907b *Ulmus protoracemosa* Penhallow: Penhallow, p. 450.
 1908 *Ulmus protoracemosa* Penhallow: Penhallow, p. 95.
 1922 *Ulmium protoracemosum* (Penhallow) Nagelhard, p. 20.
 Occurrence: Oligocene; British Columbia (Kettle River).
 Pleistocene; New York (Elmira).

Ulmium simrothi (Platen).

- 1908 *Ulmoxylon simrothi* Platen, p. 26, pl. I, f. 5, 6.
 Occurrence: Pliocene; California (Calistoga).

? **Ulmium** sp.

- 1888 *Ulmus*: Dawson, p. 33.
 Remarks: No description or figure. Requires confirmation.
 Occurrence: Upper Cretaceous (Fort Pierre Group); Western
 Canada (Head of Swift Current).

[Ulmoxylon Kaiser, 1879.]

Synonym of **Ulmium**, q. v. for all species described
 under **Ulmoxylon**.

See also **Ulmus** for **Ulmoxylon** sp.

Ulmus sp.

- 1926 *Ulmoxylon* sp.: Fietz, pp. 229, 236, pl. V, f. 9.
 1926a *Ulmus*: Fietz, p. 418.
 Remarks: See discussion on nomenclature in the Introduction.
 Occurrence: Pleistocene; Silesia and Czechoslovakia.

[Ungerites Schleiden, 1855, p. 37.]**[Ungerites tropicus** Schleiden.]

Synonym of **Ficoxylon tropicum**, q. v.

Vaccinium L.**Vaccinium** sp.

- 1914 *Vaccinium* sp.: Szafer, p. 349.
 Occurrence: Pleistocene; Poland.

[Vaticoxylon den Berger, 1927, p. 497.]

For woods of *Dipterocarpaceae* belonging to the *Vaticeae* and probably the *Vateriae*. No fossil species mentioned.

Viburnum L.**Viburnum ? sp.**

1926a *Viburnum* ? : Fietz, p. 420.

Remarks: Fietz draws attention to the close structural resemblance between the woods of *Viburnum* and *Cornus*.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

Viscum L.**Viscum album L.**

1926 *Viscum album* L.: Fietz, p. 233, pl. V, f. 3.

Remarks: On coniferous wood.

Occurrence: Pleistocene; Silesia.

Vitis L.**Vitis sp.**

1926a *Vitis*: Fietz, p. 419.

Occurrence: Prehistoric (Loess); Czechoslovakia (Brno).

? Vitis sp.

1920 ? *Vitis* sp.: Kräusel, p. 457, pl. XXXIV, f. 5; XXXVIII, f. 8, 9; text-fig. 10, 11.

Occurrence: Miocene; Silesia (Oppeln).

[**Vitoxylon** Schuster, 1911, p. 541.]

Synonym of **Ampeloxylon**, q. v. for species described as **Vitoxylon**.

Withamia Unger, 1842, p. 102.

Withamia stiriaca Unger.

1842 *Withamia stiriaca* Unger, p. 102.

1842a *Withamia stiriaca* Unger: Unger, p. 177.

1845 *Withamia stiriaca* Unger: Unger, p. 261.

1850 *Withamia stiriaca* Unger: Unger, p. 522.

1890 *Withamia stiriaca* Unger: Kaiser, p. 38.

Occurrence: Tertiary; Styria (Nestelbach).

[**Woburnia** Stopes, 1912, p. 91.]

Remarks: I agree with Kräusel in sinking this genus in *Dipterocarpoxyton*, just as *Kloedenia* has been sunk in *Quercinium*. Actually wood of the *Dipterocarpaceae* was first described as *Bredaea*. For *Woburnia porosa* Stopes, see *Dipterocarpoxyton porosum*.

Zittelia Felix, 1882, p. 73.

Zittelia elegans Felix.

1882 *Zittelia elegans* Felix, p. 73, pl. I, f. 2.

1883a *Zittelia elegans* Felix: Felix, p. 14, pl. II, f. 1, 2.

1890 *Zittelia elegans* Felix: Kaiser, p. 33.

Remarks: Relationship uncertain, perhaps *Leguminosae*.

Occurrence: Tertiary; Antigua.

Systematic List of Genera.

The families are arranged alphabetically. The numbers in brackets indicate the number of named species in each genus. Doubtful genera or species are sometimes mentioned under more than one family. Synonyms and nomina nuda are omitted entirely. Pleistocene woods and woods not specifically named are omitted from the total number of species. The attributions to families are by no means certain.

- Aceraceae: *Acerinium* (5), *Acer* sp.
- Anacardiaceae: *Anacardioxylon* (2), *Rhodium* (3).
- Araliaceae: *Aralinium* (4).
- Betulaceae: *Alnoxylon* (3), *Betulinium* (11), *Carpinoxylon* (3), *Alnus* sp., *Betula* sp., *Carpinus* sp., *Corylus* sp.
- Burseraceae: *Sumatroxylon* ? (1).
- Capparidaceae: *Capparidoxylon* (1).
- Caprifoliaceae: *Viburnum* sp.
- Casuarinaceae: *Casuaroxylon* (2).
- Celastraceae: *Celastrinoxylon* (1), *Elaeodendroxylon* (1), *Euonyminium* (1), *Rohlfisia* (1).
- Combretaceae: *Combretacinium* (1).
- Cornaceae: *Cornoxylon* (7).
- Dipterocarpaceae: *Dipterocarpoxylon* (11).
- Ebenaceae: *Ebenoxylon* (6).
- Ericaceae: *Dryoxylon sambiense* ?, *Vaccinium* sp.
- Euphorbiaceae: *Euphorbioxylon* (1), *Paraphyllanthoxylon* (1).
- Fagaceae: *Fegonium* (5), *Nothofagoxylon* (1), *Quercinium* (25), *Fagus* sp., *Castanopsis* sp., *Quercus* sp.
- Hamamelidaceae: *Hamamelidoxylon* (1), *Liquidambaroxylon* (2).
- Hippocrateaceae: *Hippocrateoxylon* (1).
- Juglandaceae: *Juglandinium* (6), *Mirbellites* ? (2).
- Lauraceae: *Laurinium* (27), *Cinnamomum* sp., *Tetranthera alnoides* ?.
- Leguminosae: *Acacioxylon* (3), *Caesalpinioxylon* (4), *Cassioxylon* (3), *Cercidoxylon* (1), *Felixia* (1), *Fichtelites* ? (1), *Hauera americana* ?, *Mohlites* ? (2), *Palackya* (1), *Pterocarpus santalinus*, *Taenioxylon* (8), *Schleidenites* (1), *Zittelia* ? (1).

Loranthaceae:	<i>Viscum album</i> .
Magnoliaceae:	<i>Dryoxylon laxum</i> ?, <i>D. silvaticum</i> ?, <i>Rosthornia</i> ? (1).
Meliaceae:	<i>Swietenioxylon</i> (1).
Menispermaceae:	<i>Lillia</i> (1).
Moraceae:	<i>Ficoxylon</i> (4), <i>Ficus callosa</i> ?.
Myrtaceae:	<i>Eucalyptus</i> sp., <i>Eugenia cordata</i> .
Oleaceae:	<i>Ornoxylon</i> (1), <i>Fraxinus</i> sp.
Piperaceae:	? <i>Piper</i> sp., ? <i>Nicolia moresneti</i> .
Platanaceae:	<i>Plataninium</i> (13).
Proteaceae:	<i>Constantinium</i> ? (1), <i>Banksia</i> sp.
Rhamnaceae:	<i>Reynosia</i> (1), <i>Rhamnacinium</i> (6).
Rosaceae:	<i>Parinarioxylon</i> (1), <i>Pruninium</i> (1).
Rutaceae:	<i>Hauera americana</i> ?, <i>Klippsteinia</i> ? (1), <i>Sjögrenia</i> ? (1).
Sabiaceae:	<i>Sabiocaulis</i> (1).
Salicaceae:	<i>Salicinium</i> (7), <i>Populocaulis</i> ? (1), <i>Salix</i> sp., <i>Populus</i> sp., <i>Rosthornia</i> ? (1).
Santalaceae:	<i>Santalum</i> sp.
Sapindaceae:	<i>Djambioxylon</i> (1), <i>Fraasia</i> ? (1), <i>Sapindoxylon</i> (2), <i>Schmiedeliopsis</i> (1).
Sapotaceae:	<i>Sapotoxylon</i> (2), ? <i>Taenioxylon porosum</i> .
Saururaceae:	<i>Saururopsis</i> (1).
Simarubaceae:	<i>Simarubinium</i> (2).
Staphyleaceae:	<i>Staphylea</i> sp.
Sterculiaceae:	<i>Dombeyoxylon</i> (3), <i>Nicolia</i> (5), <i>Sillimannia</i> (1), <i>Staubia</i> (1), <i>Tarrietioxylon</i> (1), <i>Revesia wallichi</i> ?.
Ternstroemiaceae:	<i>Aptiana</i> (1), <i>Ruyschioxylon</i> (1), <i>Ternstroemiacinium</i> (2), <i>Helictioxylon schenki</i> ?.
Thymelaeaceae:	<i>Hauera americana</i> ?.
Tiliaceae:	<i>Tilioxylon</i> sp.
Ulmaceae:	<i>Ulmimum</i> (9), <i>Ulmus</i> sp.
Vitaceae:	<i>Ampeloxylon</i> (3), <i>Vitis</i> sp.

Incertae Sedis (Lianes): *Anomaloxylon* (1), *Helictoxylon* (7).

Incertae Sedis: *Aachenoxylon* (1), *Brongniartites* (1), *Bronnites* (4), *Cantia* (1), *Cottaites* (2), *Dryoxylon* (5), *Hauera* (3), *Hythia* (1), *Jugloxylon* (1), *Klippsteinia* (1), *Meyenites* (1), *Pataloxylon* (2), *Petzholdia* (3), *Piccolominites* (1), *Pritchardia* (1), *Sabulia* (1), *Stephanoxylon* (1), *Suevioxylon* (1), *Tchihatcheffites* (1), *Withamia* (1).

Total number of Generic Names (not including synonyms nor recent genera recorded from Pleistocene beds) 95.

Total number of species (not including synonyms, nomina nuda, and Pleistocene woods) 280.

Geographical Index.

Synonyms, nomina nuda, dubious and unnamed woods have been omitted, and also some Pleistocene records of recent woods.

Europe.

Germany.

Acerinium borussicum, *A. terrae-coeruleae*, *Alnoxylon inclusum*, *Ampeloxylon ampelopsoides*, *A. coheni*, *Betulinium geinitzii*, *B. lignitum*, *B. oligocaenicum*, *B. tenerum*, *B. sp.*, *Carpinoxylon compactum*, *C. sp.*, *Castanopsis sp.*, *Cornoxyylon* (6 spp.), *Dryoxylon laxum*, *D. sambiensis*, *Ebenoxylon tenax*, *Fegonium dryandrae-forme*, *F. lignitum*, *F. schenki*, *Hauera bornensis*, *Helictoxylon roemeri*, *H. speciosum*, *Juglandinium longiradiatum*, *J. triebelii*, *J. sp.*, *Laurinium algovicum*, *L. biseriatum*, *L. brunsvicense*, *L. haasi*, *L. perseoides*, *L. radiatum*, *L. tigurinum*, *L. triseriatum*, *L. sp.*, *Plataninium boreale*, *P. klebsii*, *P. megapolitanum*, *P. subaffine*, *Quercinium quercoides*?, *Q. sabulosum*, *Q. subgarryanum*, *Q. sp.*, *Rhamnacinium primaevum*, *Salicinium miocenicum*, *Suevioxylon*, *Swietenioxylon*, *Taenioxylon varians*, *T. sp.*, *Ternstroemiacinium krauseli*.

Holland.

Cornoxyylon latiporosum, *C. sp.*, *Laurinium nectandrioides*, *Salicinium varians*.

Belgium.

Aachenoxylon, *Nicolia moresneti*, *Salicinium bruxellense*.

England.

Aptiana, *Cantia*, *Dipterocarpoxyylon porosum*, *Hythia*, *Piper sp.*, *Sabulia*.

France.

Ampeloxylon cineritarum, *Betulinium parisiense*, *Hamamelidoxyylon*, *Liquidambaroxylon lecointreae*, *Quercinium eocenicum*, *Salicinium biradiatum*.

Italy.

Acerinium astianum, *Alnoxylo* sp., *Anomaloxylon*, *Betulinium paronae*, *Juglandinium mediterraneum*, *J. zuriense*, *Laurinium xyloides*, L. sp., *Piccolominites*, *Quercinium astianum*, *Salicinium antiquum*, *S. messinianum*, *Ulmium lovisatoi*, *U. pliocenicum*.

Austria.

Acerinium danubiale, *Betulinium tenerum*, *Bronnites vindobonensis*, *Cottaites vasculosus*, *Fichtelites*, *Hauera stiriaca*, *Meyenites*, *Mohlites parenchymatosus*, *Plataninium vasculosum*, *Quercinium austriacum*, *Q. sabulosum*, *Rosthornia*, *Salicinium populinum*, *Sapotoxylon*, *Ulmium lapidarium*, *Withamia*.

Hungary.

Alnoxylo vasculosum, *Betulinium priscum*, *Carpinoxylo vasculosum*, *Cassioxylo zirkeli*, *Cottaites robustior*, *Dryoxylo silvaticum*, *Fraasia*, *Helictoxylo anomalum*, *Juglandinium mediterraneum*, *J. schenki*, *Laurinium antiquum*, *L. aromaticum*, *Lillia*, *Liquidambaroxylo speciosum*, *Mohlites cribrosus*, *Plataninium porosum*, *P. regulare*, *Quercinium böckhianum*, *Q. compactum*, *Q. helictoxyloides*, *Q. leptotichum*, *Q. longiradiatum*, *Q. quercoides*, *Q. sabulosum*, *Q. vasculosum*, *Rhodium juglandinum*, *Schleidenites*, *Staubia*, *Taenioxylo pannonicum*, *Tilioxylo* sp., *Ulmium hungaricum*, *U. kleinii*.

Poland.

Betulinium parisiense ?, *B. sp.*, *Dryoxylo nivium*, *Helictoxylo roemeri*, *Ornoxylo*, *Petzholdia polonica*, *Plataninium salinarum*, *P. vasculosum*, *Quercinium transiens*, *Stephanoxylo*.

Czechoslovakia.

Betulinium stagnigenum, *Ficoxylo tropicum*, *Laurinium diluviale*, *Quercinium sabulosum*.

Roumania.

Bronnites transylvanicus, *Quercinium transylvanicum*.

Turkey.

Bronnites orientalis, *Constantinium*, *Tchihatcheffites*.

Greece.

Brongniartites, *Ebenoxylo* sp., *Juglandinium mediterraneum*, *Mirbellites lesbius*.

Russia.

Betulinium rossicum, *Euonyminium auerbachi*, *Quercinium montanum*, *Q. rossicum*, *Rhodium ungeri*.

Azerbaijan.

Anacardioxylo uniradiatum, *Combretacinium*, *Fegonium caucasicum*, *Plataninium porosum*, *P. sp.*, *Rhamnacinium affine*, *Sjögrenia*, *Taenioxylo porosum*, *Ternstroemiacinium euryoides*.

Asia.**Japan.**

Alnoxyton yezoense, Betulinium macclintockii, Fegonium hokkaidense, Jugloxyton, Populocaulis, Sabiocaulis, Saururopsis.

Burma.

Dipterocarpoxyton burmense.

Annam.

Dipterocarpoxyton annamense.

Philippines.

Helictoxyton luzonense, Palackya, Rhodium philippinense, Tainioxyton eperuoides.

East Indies.

Dipterocarpoxyton swedenborgii, Pterocarpus santalinus, Tainioxyton blanfordi, T. pacificum.

Sumatra.

Caesalpinioxyton Palembangense, Dipterocarpoxyton djambiense, D. kräuseli, D. tobleri, Djambioxyton, Ruyschioxyton, Sapindoxyton janssoni, Sumatroxyton, Tarrietioxyton.

Java.

Dipterocarpoxyton goepperti, D. javanense, D. moroides, D. spectabile, Helictoxyton schenki, Hippocrateoxyton javanicum, Parinarioxyton, Sapindoxyton klitzingi.

Australasia.**Australia.**

Banksia sp., Caesalpinioxyton oweni, Eucalyptus sp., Pataloxyton (2 spp.).

New Guinea.

Laurinium meyeri.

New Caledonia.

Nicolia caledonica.

Bismarck Archipelago.

Caesalpinioxyton oweni.

New Zealand.

Nicolia zelandica.

Africa.**Rio d'Oro.**

Caesalpinioxylon quirogae, Nicolia aegyptiaca.

Algeria.

Cassioxylon bartholomei.

Tunisia.

Acacioxylon antiquum, Ebenoxylon tunetanum, Ficoxylon cretaceum, Nicolia aegyptiaca.

Libyan Desert (see also Egypt).

Laurinium desioi, Nicolia aegyptiaca, N. giarabubensis, Rohlfisia.

Sirtica.

Ficoxylon cretaceum, Laurinium desioi.

Egypt.

Acacioxylon antiquum, A. vegae, Acerinium aegyptiacum, Caesalpinioxylon oweni, Capparidoxylon geinitzi, Celastrinoxylon affine, Dombeyoxylon aegyptiacum, Ebenoxylon ebenoides, Ficoxylon cretaceum, Laurinium primigenium, Nicolia aegyptiaca.

Sudan.

Caesalpinioxylon oweni.

Abyssinia.

Dombeyoxylon affine, Nicolia aegyptiaca.

Somaliland.

Nicolia aegyptiaca.

Madagascar.

Laurinium albiense.

South Africa.

Eugenia cordata.

North America.**Banksland.**

Betulinium macclintockii.

Canada.

Plataninium sp., Uminium columbianum, U. protoamericanum, U. proracemosum.

United States.

Anacardioxylon magniporosum, Aralinium (4 spp.) Carpinoxylon pfefferi, Cercidoxylon zirkeli, Dombeyoxylon jacksonensis, Dryoxylon americanum, Ebenoxylon boreale, E. speciosum, Elaeodendroxylon, Felixia, Ficoxylon, helictoxyloides, Helictoxylon wilcoxianum, Laurinium aromaticum, L. bakeri, L. brandonianum, L. branneri, L. californicum, L. eberi, L. lesquerianum, L. pulchrum, L. wilcoxianum, L. sp., Paraphyllanthoxylon, Plataninium crystallophorum, P. haydeni, P. knowltoni, P. pacificum, Pruninium, Quercinium abromeiti, Q. anomalum, Q. knowltoni, Q. lamarense, Q. lesquereuxi, Q. pliocaenicum, Q. solerederi, Q. wardi, Q. sp., Reynosia texana, Rhamnacinium porcupinianum, R. radiatum, R. texanum, R. triseriatum, Sillimannia, Simarubinium, Ulminium simrothi.

Mexico.

Acacioxylon tenax, Hauera americana.

Guatemala.

Laurinium guatemalense.

Central America.

Taenioxylon multiradiatum.

Antigua.

Anacardioxylon spondiaeforme, Bronnites antiguensis, Cassioxylon anomalum, Ebenoxylon diospyroides, Hauera americana, Helictoxylon tenerum, Petzholdia tropica, Schmiedeliopsis, Taenioxylon irregulare, T. multiradiatum, T. varians, Zittelia.

West Indies.

Petzholdia major, Pritchardia.

South America.**Columbia.**

Euphorbioxylon.

Brazil.

Taenioxylon ingaeforme.

Juan Fernandez.


Santalum sp.

Uruguay.

Caesalpinioxylon nathorstii

Argentina.

Betulinium rocae.



Patagonia.

Laurinium uniseriatum, *Nothofagoxylon scalariforme*.

Chile.

Nothofagoxylon sp.

Tierra del Fuego.

Nothofagoxylon sp.

Antarctica.

Laurinium uniseriatum, *Nothofagoxylon scalariforme*.

Addenda.

Since the completion of this work I have seen the following papers:

Chiarugi, A. 1929. La presenza in Sardegna di elementi paleoxilologici sahariani. *Nuovo Giorn. Bot. Ital.*, XXXVI, pp. 254—258.

Chiarugi, A. 1930. Contribuzione alla paleoxilologia dell'Africa. *Compte Rendu XV Int. Geol. Congress, Pretoria, II*, pp. 179—182, 1 map.

The first paper records the presence in the Miocene of Sardinia of *Dombeyoxylon aegyptiacum* and *Laurinium desioi*, and suggests that *Ulmium lovisatoi* Falqui is probably identical with the *Dombeyoxylon*. The second paper discusses geographical distribution and records the following (all *nomina nuda*) from Somaliland: *Dipterocarpoxylon somalense*, *D. giubense*, *D. scebelianum*, *Sapindoxylon benadirensis*.



Index.

An asterisk denotes a living species.

- Aachenosaurus multidentis 16, 57
 Aachenoxylon 4, 16
 abromeiti, Quercinium 63
 Acacioxylon 16
 Acer 16, 17
 Acerinium 17
 acerinum, Plataninium 60
 aceroides, Plataninium 60, 62
 aegyptiaca, Nicolia 31, 55, 57
 aegyptiacum, Acerinium 17
 aegyptiacum, Dombeyoxylon 36,
 90
 aequimontanus, Meyenites 54
 affine, Celastrinoxylon 29
 affine, Dombeyoxylon 37
 affine, Rhamnacinium 69, 70
 alata, Cassia 28
 albiense, Laurinium 48
 album, Viscum 21, 82
 algovicum, Laurinium 48
 alnoides, Tetranchera 79
 Alnoxylon 17, 18
 Alnus 18
 Ambaroxylon 18, 54
 americana, Hauera 44
 americanum, Dryoxylon 37
 Ampelopsis 18
 ampelopsoides, Ampeloxylon 18
 Ampeloxylon 18
 Anacardiaceae 69, 77
 Anacardioxylon 19, 76
 anglia, Casuaroxylon 28
 Anisoptera 34
 annamense, Dipterocarpoxyton 34
 Anomaloxylon 19
 anomalum, Cassioxylon 28
 anomalum, Helictoxylon 44
 anomalum, Quercinium 63
 Anonaceae 74
 antiguensis, Bronnites 25
 antiquum, Acacioxylon 16
 antiquum, Laurinium 49
 antiquum, Salicinium 72
 Aptiana 19
 Aralinium 20
 arborescens, Cantia 27
 arizonense, Paraphyllanthoxylon
 58
 aromaticum, Laurinium 49
 articulatus, Fichtelites 42
 astianum, Acerinium 17
 astianum, Quercinium 64
 Aquifoliaceae 20
 Aquilarineae 44
 auerbachi, Euonyminium 40
 Aulacoxylon 20
 Aurantiaceae 44, 48, 76
 australe, Banksioxylon 20
 austriacum, Quercinium 64
 bakeri, Laurinium 49, 52
 Banksia 20, 32
 Banksioxylon 20
 bartholomei, Cassioxylon 28
 benadirensis, Sapindoxylon 90
 Betula 21, 24, 32
 Betulaceae 18, 33
 Betulinium 21—24, 33, 50
 Betuloxylon 21, 24, 32
 biradiatum, Salicinium 72
 biseriatum, Laurinium 49, 52
 blanfordi, Taenioxylon 77
 böckhianum, Quercinium 64, 65
 bohemicum, Ficoxylon 42
 Bombacaceae 33
 bornensis, Hauera 44
 boreale, Ebenoxylon 38
 boreale, Plataninium 60
 borussicum, Acerinium 17
 brandonianum, Laurinium 49
 branneri, Laurinium 49
 Bredaea 25, 35, 82
 Bridelia 58
 Brongniartites 4, 25
 Bronnites 4, 25
 brunswicense, Laurinium 50
 bruxellense, Salicinium 72

- burmense, Dipterocarpoxyton 34
 Burseraceae 77
 byzantina, Tchihatcheffites 78

 *Caesalpinia echinata 26
 Caesalpinaceae 44, 58, 77.
 Caesalpinioxyton 25
 Caesalpinium 26
 caledonica, Nicolia 56
 californicum, Laurinium 50
 callosa, Ficus 43
 camphora, Cinnamomum 29
 Cantia 4, 27
 Capparidoxyton 27
 Capparis 27
 Caprifoliaceae 20, 33
 carinthiaca, Rosthornia 71
 Carpinoxyton 27, 52
 Carpinus 27
 Carya 47
 Cassia 28
 *Cassia speciosa 28
 Cassioxyton 28
 Castanopsis 28
 Casuaroxyton 28
 cathartica, Rhamnus 70
 caucasicum, Fegonium 41
 Cedrelaceae 32
 Celastraceae 29, 48, 71
 Celastrinoxyton 29
 celastroides, Rohlfsia 71
 Celtites (Celtitis) 29, 80
 Cercidoxyton 29
 cerris, Quercus 68
 Charpentieria 29, 38
 cineritarum, Ampeloxylon 18
 Cinnamomum 29, 49
 coheni, Ampeloxylon 18
 columbianum, Ulminium 80
 Combretacinium 29
 compactum, Carpinoxyton 27
 compactum, Quercinium 64, 65
 compositus, Schleidenites 74
 Constantinium 29
 cordata, Eugenia 39
 Cornoxyton 29—31
 Cornus 29, 30, 82
 *Cornus alba 30
 Corylus 31
 *Coscinium fenestratum 54
 Cottaites 4, 31, 80
 cretaceum, Cornoxyton 29
 cretaceum, Ficoxyton 42
 cribrusos, Mohlites 55
 crystallophilum, Plataninium 60
 crystallophora, Sjögrenia 76
 crystallophorum, Simarubinium 75
 Cyrillaceae 20

 danubiale, Acerinium 17
 danubiense, Acerinium 17
 desioi, Laurinium 50, 90
 Dicotyledonous wood 31—34, 44,
 59
 *Dicypellium caryophyllatum 49
 diluviale, Betulinium 24
 diluviale, Laurinium 24, 50
 diospyroides, Ebenoxyton 38
 Diospyros 34
 *Diospyros discolor 38
 *Diospyros ebenus 39
 *Diospyros virginiana 38
 Dipterocarpaceae 26, 34—36, 55,
 81, 82
 Dipterocarpoxyton 34, 37, 75, 82,
 90
 Dipterocarpus 34
 djambiense, Dipterocarpoxyton 34
 Djambioxyton 36
 *Dombeya mollis 37
 Dombeyoxyton 36, 76, 90.
 dryandraeforme, Fegonium 41
 Dryobalanops 37
 Dryobalanoxylon 35—37
 Dryoxyton 4, 37, 38, 77
 dubium, Stephanoxylon 76

 Ebenaceae 38
 ebenoides, Ebenoxyton 38, 39
 Ebenoxyton 34, 38
 eberi, Laurinium 50
 Elaeodendroxylon 39
 elegans, Miquelites 54
 elegans, Zittelia 82
 elgari, Hythia 46
 engelhardti, Simarubinium 76
 eocenicum, Quercinium 64
 *Eperua decandra 77
 eperuoides, Taenioxyton 77
 Erica 37
 *Erica vagans 38
 Ericaceae 33
 eriodendroides, Staubia 76
 Eriodendron 36, 76
 erraticum, Cornoxyton 30
 Erythrobalanus 65
 Eucalyptus 39
 Eugenia 39
 Euonymus 40
 *Euonymus japonicus 40
 Euonyminium 40
 Euphorbia 40
 Euphorbiaceae 40, 58
 Euphorbioxyton 40, 58
 *Eurya acuminata 20
 *Eurya latifolia 79
 euryoides, Ternstroemiacinium 79
 excellens, Aralimum 20
 excelsior, Fraxinus 43
 Exogenites 4, 22, 37, 40

- extorre, Casuaroxylon 28
 Fagoxylon 40
 Fagus 40, 46
 Fegonium 40, 41, 61, 62
 Felixia 41
 Fichtelia 42
 Fichtelites 4, 42
 Ficoxylon 42
 Ficus 43
 *Ficus cordata 42
 *Ficus elastica 43
 *Ficus sycomorus 42
 Fraasia 4, 43
 fraxinoides, Ornoxylon 38
 Fraxinoxylon 43
 Fraxinus 43, 58
 *Fraxinus excelsior 43

 geinitzi, Capparidoxylon 27
 geinitzii, Betulinium 21
 giarabubensis, Nicolia 57
 giubense, Dipterocarpoxyton 90
 goepperti, Dipterocarpoxyton 34
 graecus, Brongniartites 25
 Grewia 35
 Grewioxylon 35, 44
 guatemalense, Laurinium 4, 51
 Guazuma 36
 gumbelii, Sapotoxylon 74
 gunmosum, Pruninium 63

 haasi, Laurinium 51
 Hamamelidoxylon 44
 Hamamelis 31
 hamoanum, Jugloxylon 47
 hasskarlianus, Piperites 60
 Hauera 44
 haydeni, Plataninium 60
 helictoxyloides, Ficoxylon 42
 helictoxyloides, Quercinium 64
 Helictoxyton 19, 44, 71
 *Hippocratea scandens 46
 *Hippocratea viridis 46
 Hippocrateoxylon 45
 Hippomane 40
 hokkaidense, Fegonium 41
 holsatae, Cornoxylon 30
 holsaticum, Cornoxylon 30
 Hopea 34
 Humiriaceae 33
 hungaricum, Ulminium 80
 Hythia 4, 46

 inclusum, Alnoxylon 17
 *Inga fastuosa 77
 ingaeforme, Taenioxylon 77
 insignis, Pritchardia 63
 irregulare, Taenioxylon 77
 itersonii, Parinarioxylon 58

 jacksonensis, Dombeyoxylon 37
 janssonii, Sapindoxylon 73
 Jatropha 40
 javanense, Dipterocarpoxyton 35
 javanicum, Hippocrateoxylon 45
 javanicus, Junghuhnites 48
 jenense, Dryoxylon 37
 Jordania 38, 39, 46
 Juglandaceae 55
 Juglandinium 4, 46, 55
 juglandinum, Rhoidium 4, 70
 Juglandoxylon 46, 47
 Juglans 46, 47
 *Juglans australis 47
 Juglansoxylon 47
 Jugloxylon 4, 47
 Junghuhnites 48

 klebsii, Plataninium 60
 kleinii, Ulminium 80
 Klippsteinia 4, 48
 klitzingi, Sapindoxylon 74
 Kloedenia 48, 63, 66
 knowltoni, Plataninium 61
 knowltoni, Quercinium 64, 65
 kräuseli, Dipterocarpoxyton 35
 kräuseli, Ternstroemiacinium 79

 lamarense, Quercinium 65
 lapidariorum, Ulminium 80
 latiporosum, Cornoxylon 30
 latiradiata, Felixia 41
 Lauraceae 33
 Laurinium 4, 48—53, 90
 Laurinoxylon 48—53
 Lauroxylon 53
 Laurus 51—53
 *Laurus benzoin 53
 *Laurus nobilis 50, 52
 *Laurus obtusifolia 49
 *Laurus sassafras 49
 laxum, Dryoxylon 38
 lecointreae, Liquidambaroxylon 54
 Leguminoseae 31, 33, 42, 55, 68,
 75, 80, 82
 leptotichum Quercinium 65
 lesbius, Mirbellites 46, 55
 lesquereuxi, Quercinium 65
 lesquerianum, Laurinium 51
 lignitum, Betulinium 21, 24
 lignitum, Fegonium 41
 Lillia 53
 lindgreni, Araliniium 20
 Liquidambar 31
 *Liquidambar styraciflua 54
 Liquidambaroxylon 54
 longiradiatum, Juglandinium 46
 longiradiatum, Quercinium 65
 lovisatoi, Ulminium 80, 90
 luzonense, Helictoxyton 45

- macclintockii, Betulinium 22
 magniporosum, Anacardioxylon 19
 Magnolia 38, 54
 Magnoliaceae 20, 71
 Magnolites 38, 54
 magnoradiatum, Anomaloxyton 19
 major, Petzholdia 59
 mediterraneum, Juglandinium 4, 46, 55
 medullaris, Klippsteinia 48
 megapolitanum, Plataninium 60
 Meliaceae 32, 77.
 melliodora, Eucalyptus 39
 Menispermaceae 54
 Meyenites 4, 54
 meyeri, Laurinium 51
 messinianum, Salicinium 72
 Mimoseae 77
 minor, Nicolia 26
 miocenicum, Salicinium 4, 72, 73
 Miquelites 54
 Mirbellites 4, 46, 55
 *Mithridatea erecta 52
 Mohlites 4, 55
 mollii, Sumatroxylon 76, 77
 montanum, Quercinium 65
 moresneti, Nicolia 16, 57
 moroides, Dipterocarpoxyton 35
 multident, Aachenosaurus 16, 57
 multident, Aachenoxylon 16
 multiradiatum, Aralinium 20
 multiradiatum, Taenioxyton 77
 Myrica 30
 myricaeforme, Cornoxyton 30
 Myrtaceae 33, 38

 nathorsti, Caesalpinioxyton 25
 Naucleoxyton 35
 Nectandra 49, 51
 nectandrioides, Laurinium 51
 Nicolia 26, 31, 55—57, 59
 niponensis, Saururopsis 74
 nivium, Dryoxyton 38
 Nothofagoxyton 57, 58
 *Nothofagus betuloides 57
 Nyctomyces antediluvianus 55
 Nyctomyces entoxylinus 56

 Ocotea 48, 52
 Ocoteoxyton 48, 52, 58
 Oleaceae 58
 oligocaenicum, Betulinium 21, 22
 Oreodaphne 49
 orientalis, Bronnites 25
 Ornoxyton 43, 58
 oweni, Caesalpinioxyton 26

 pacificum, Plataninium 61
 pacificum, Taenioxyton 78
 Palackya 4, 58

 palembangense, Caesalpinioxyton 26
 pannonicum, Taenioxyton 78
 Papilionaceae 77
 Paraphyllanthoxyton 58
 parenchymaticum, Aralinium 20
 parenchymatosus, Mohlites 55
 Parinarioxylon 58
 *Parinarium sumatranum 58
 parisiense, Betulinium 22
 paronae, Betulinium 22
 Parrotia 44
 Pataloxyton 4, 58
 Persea 48, 49
 *Persea gratissima 50, 51
 perseoides, Laurinium 51
 Perseoxyton 48—50, 53, 59
 Petzholdia 4, 59
 pfefferi, Carpinoxyton 27, 52
 Phegonium 40, 59, 61
 philippinense, Palackya 58
 philippinense, Rhodium 70
 Phyllanthus 58
 Piccolominites 4, 59
 Piper 59
 Piperaceae 57
 piperita, Eucalyptus 39
 Piperites 60
 Plataninium 31, 40, 60—62
 Platanus 60, 62
 *Platanus occidentalis 60
 pliocaenicum, Quercinium 65
 pliocenicum, Ulminium 80
 polonica, Petzholdia 59
 polymorphum, Elaeodendroxylon 39
 Pomaderris 69
 Pomelia 73
 populinum, Salicinium 4, 72
 Populocaulis 4, 62
 Populoxyton 62
 Populus 33, 62, 63, 69, 70, 72, 73
 porcupinianum, Rhamnacinium 69
 porosum, Dipterocarpoxyton 35
 porosum, Pataloxyton 58
 porosum, Plataninium 61
 porosum, Taenioxyton 78
 primaevum, Quercinium 66, 67
 primaevum, Rhamnacinium 69, 70
 primigenium, Laurinium 51
 Prinos 69
 priscum, Betulinium 23
 Pritchardia 4, 63
 Proteaceae 29
 proteoides, Constantinium 29
 Protocedroxylon 19
 Protocupressinoxyton 19
 protoracemosum, Ulminium 81
 Pruninium 63
 Prunus 63

- Pterocarpus 63
 *Pterocarya caucasica 47
 Pterospermum 76
 pulchrum, Laurinium 52

 Quercinium 33, 63—68
 Quercites 66, 67
 quercoides, Quercinium 66—68
 Quercoxylon 68
 Quercus 65—69
 *Quercus castanaefolia 66, 67
 *Quercus garryana 67
 *Quercus lusitanica 64
 *Quercus pedunculata 66
 *Quercus sessiliflora 65
 *Quercus tozza 65
 quirogae, Caesalpinioxylon 26
 *Quisqualis pubescens 29
 quisqualoides, Combretacinium 29

 radiata, Aptiana 19
 radiatum, Laurinium 52
 radiatum, Rhamnacinium 62, 70
 regulare, Plataninium 61
 renaulti, Hamamelidoxylon 44
 Reveesia 69
 Reynosia 69
 *Reynosia latifolia 69
 Rhamnacinium 4, 62, 69, 70, 73, 80
 Rhamnus 70
 Rhizoalnoxylon 17, 70
 Rhodium 4, 70
 *Rhus juglandifolia 70
 robustior, Cottaites 31
 rocae, Betulinium 23, 33
 roemeri, Helictoxylon 45
 Roemeria 37, 70
 Rohlfisia 29, 71
 Rosaceae 58
 Rosthonia 4, 71
 rossicum, Betulinium 23
 rossicum, Quercinium 65, 66
 Rubiaceae 52
 Ruizia 36
 Ruyschia 71
 Ruyschioxylon 45, 71

 *Sabia japonica 71
 Sabiocalis 71
 Sabulia 4, 71
 sabulosum, Quercinium 66
 sakuraii, Sabiocalis 71
 Salicaceae 71—73
 Salicinium 4, 62, 72
 Salicinoxylon 4, 72, 73
 salinarum, Plataninium 61
 Salix 4, 37, 62, 72, 73
 salzhauseusis, Betula 21, 24
 sambiense, Dryoxylon 38
 sambiensis, Erica 37

 santalinus, Pterocarpus 63
 Santalum 73
 *Santalum album 73
 Sapindaceae 36, 43, 74, 75
 sapindoides, Fraasia 43
 Sapindopsis 74
 Sapindopsoxylon 73
 Sapindoxylon 73, 74, 90
 Sapotaceae 74
 Sapotoxylon 74, 78
 sardus, Piccolominites 59
 Saururaceae 74
 Saururopsis 74
 Saxifragaceae 33
 scalariforme, Nothofagoxylon 57
 scalariforme, Pataloxylon 59
 scebelianum, Dipterocarpoxyton 90
 schenki, Fegonium 41
 schenki, Ficoxylon 42
 schenki, Helictoxylon 45, 71
 schenki, Juglandinium 46
 Schimperites 63, 65, 74
 Schinus 4, 69
 *Schinus molle 69
 Schleidenites 4, 74
 Schmidites 68, 75
 *Schmiedelia haemorrhoea 75
 Schmiedeliopsis 75
 schuchii, Mirbellites 55
 scottii, Sabulia 71
 scottii, Woburnia 35
 Shorea 34
 Shoreae 75
 Shoreoxylon 26, 34—36, 75
 silvaticum, Dryoxylon 38
 Simarubaceae 19
 Simarubinium 75
 simrothi, Ulminium 81
 Sillimannia 4, 75
 Sjögrenia 4, 76
 solerederi, Quercinium 67
 solidior, Cornoxylon 30
 somalense, Dipterocarpoxyton 90
 sparnacense, Aulacoxylon 20
 spectabile, Dipterocarpoxyton 35
 speciosum, Ebenoxylon 39
 speciosum, Euphorbioxylon 40
 speciosum, Helictoxylon 45
 speciosum, Liquidambaroxylon 54
 spondiaeforme, Anacardioxylon 19
 *Spondias lutea 19
 stagnigenum, Betulinium 23
 Staphylea 76
 staubi, Quercinium 65, 67
 Staubia 76
 Stephanoxylon 4, 76
 Sterculiaceae 26, 36, 37, 75, 78
 sternbergense, Swietenioxylon 77
 stiriaca, Hauera 44
 stiriaca, Withamia 82



- subaffine, Plataninium 61
 subgarryanum, Quercinium 66, 67
 Suevioxylon 4, 76
 sumatrense, Djambioxylon 36
 sumatrense, Ruyschioxylon 71
 sumatrense, Tarrietioxylon 78
 Sumatroxylon 4, 76
 swedenborgii, Dipterocarpoxyton
 35
 *Swietenia mahagoni 77
 Swietenioxylon 77

 taeniatum, Sapotoxylon 74
 Taenioxylon 77
 Tarrietia 78
 Tarrietioxylon 78
 tasmanicum, Banksioxylon 21
 Tehihatcheffites 4, 78
 Tehihatchewia 78
 tenax, Acacioxylon 16
 tenax, Ebenoxylon 34, 39
 tenerum, Betulinium 23
 tenerum, Helictoxylon 45
 Ternstroemiaceae 20, 45, 71
 Ternstroemiacinium 79
 Ternstroemioxylon 79
 terrae-coeruleae, Acerinium 17
 Tetrathera 79
 texana, Reynosia 69
 texana, Sillimannia 75
 texanum, Rhamnacinium 70
 Tilia 79
 Tilioxylon 79
 tigurinum, Laurinium 52
 tirolense, Caesalpinium 26
 tobleri, Dipterocarpoxyton 35, 36
 transiens, Quercinium 67
 transylvanicus, Bronnites 25
 transylvanicum, Quercinium 67
 triebelii, Juglandinium 47
 triseriatum, Laurinium 52
 triseriatum, Rhamnacinium 70
 Trochodendromagnolia 4, 79
 tropica, Petzholdia 59
 tropicum, Ficoxylon 42
 tunetana, Nicolia 57
 tunetanum, Ebenoxylon 39

 Ulmaceae 80
 Ulminium 50, 80, 81
 Ulmoxyton 80, 81
 Ulmus 80, 81

 *Ulmus americana 81
 ungeri, Rhoidium 70
 Ungerites 42, 81
 uniradiatum, Anacardioxylon 19
 uniseriatum, Laurinium 52

 Vaccinium 20, 81
 varians, Salicinium 72
 varians, Taenioxylon 78
 vasculosum, Alnoxyton 18
 vasculosum, Carpinoxylon 27
 vasculosum, Plataninium 61
 vasculosum, Quercinium 66, 68
 vasculosus, Cottaites 31
 vateri, Cornoxylon 30
 Vateriae 81
 Vaticaeae 81
 Vaticoxylon 81
 vegae, Acacioxylon 16
 Viburnum 82
 vicentinum, Anomaloxylon 19
 viennensis, Bronnites 25
 vindobonensis, Bronnites 25
 Viscum 21, 82
 Vitaceae 18
 viticulosa, Lillia 53
 Vitis 18, 82
 Vitoxylon 18, 82

 wallichi Reveesia 69
 wardi, Quercinium 68
 wichmanni, Juglandinium 47
 wiedemanni, Nicolia 56
 wilcoxianum, Helictoxylon 45
 wilcoxianum, Laurinium 52
 Withamia 4, 82
 Woburnia 35, 82

 xyloides, Laurinium 4, 53

 yezoëense, Alnoxyton 18
 yezoënsis, Populocaulis 62

 zelandica, Nicolia 57
 zirkeli, Cassioxylon 28
 zirkeli, Cercidoxylon 29
 zirkeli, Ficoxylon 43
 zirkeli, Schmiedeliopsis 75
 Zittelia 4, 82
 zonatum, Suevioxylon 76
 zuriense, Juglandinium 47
 Zygomphyllaceae 54



- Pars 42: Ch. Schuchert et Clara M. LeVene, Brachiopoda. 1929. 140 p.
44: J. Felix, Anthozoa pliocenica et pliocenica [cum indice
ad omnes partes Anthozoarum tertiarium et quaternarium].
45: F. Plieninger, Pterosauria. 1930. 84 p.
47: G. G. Simpson, Post-Mesozoic Marsupialia. 1930. 87 p.
48: V. van Straelen, Crustacea Eumalacostraca (Decapodis exclu-
sis). 1931. 98 p.

In Vorbereitung:

- K. Andrée, Insecta et Vertebrata succini.
W. O. Dietrich, Orbitolinidae.
K. Freudentberg, Palaeocrinoidea.
W. Freudentberg, Mammalia fossilia: Hominidae.
H. Gerth, Tetracoralla carbonica et permiana.
Porifera palaeozoica.
Haber, Gastropoda liassica.
A. Handlirsch, Insecta fossilia II: Mesozoica.
Insecta fossilia III: Caenozoica.
K. Hummel, Trionychia fossilia.
W. Janensch, Saurischia [Theropoda et Sauropoda] jurassica et
cretacea.
E. Jaworski, Lamellibranchiata mesozoica.
A. Kieslinger, Nautiloidea.
A. Liebus, Foraminifera.
H. Matern, Ostracoda palaeozoica.
H. Rauff, Porifera (palaeozoicis exclusis).
Ch. Resser, Trilobitae cambricae.
R. Richter, Trilobitae devonicae et carbonicae. II, Vermes, Phyllo-
poda, Xiphosura.
F. A. Schilder, Cypraeidae, Triviidae et Amphiperasidae (=
Ovulidae) fossiles.
H. Scupin, Spiriferacea liassica.
J. Smith, Cephalopoda carbonica.
E. Stolley, Ammonoidea infracretacea.
V. van Straelen, Phyllocarida.
W. Teppner, Lamellibranchiata tertiaria. 'Anisomyaria'. III: Spon-
dylidae, Anomiidae, Ostreidae, Mytilidae.
E. Voigt, Bryozoa cheilostomata cretacea.
R. Wedekind, Tetracoralla silurica et devonica.

Fortsetzung siehe 4. Umschlagseite.

Fossilium Catalogus

II: Plantae:

- Pars 1, 15, 16: W. Jongmans, Lycopodiales. (incl. Hydropterideae, Psilophytales, Sphenophyllales). I—III. 1913—1930. 860 p.
2—5, 7, 9, 11: W. Jongmans, Equisetales. 1914—1924. 7 partes. 831 p.
6: K. Nagel, Juglandaceae. 1915. 87 p.
8: K. Nagel, Betulaceae. 1916. 177 p.
10: K. Nagalhard [vormals Nagel], Ulmaceae. 1922. 86 p.
12: O. Posthumus, Inversicatenales (Botryopterideae et Zygopterideae). 1926. 56 p.
13: H. N. Dixon, Muscineae. 1927. 116 p.
14: W. N. Edwards et F. M. Wonnacott, Sapindaceae. 1928. 84 p.

In Vorbereitung:

- W. N. Edwards, Angiospermae [pro parte]. Anacardiaceae; weiter: Aceraceae, Celastraceae, Aquifoliaceae, Buxaceae, Coriariaceae, Hippocastaneae, Icacinaceae, Staphyleaceae, Melianthaceae, Sabiaceae; Magnoliaceae, Sterculiaceae, Ebenaceae, Sapotaceae, Symplocaceae, Styracaceae.
R. Florin, Coniferae.
J. Groves, Charaphytae.
L. Hoerhammer, Monokotylen.
W. Jongmans, Filices, Semina palaeozoica, Lycopodiales IV.
R. Kräusel, Plantae devonicae.
L. Laurent, Angiospermae [pro parte].
H. Peragallo, Diatomeae.
J. Pia, Algae.
J. Schuster, Cycadophyta.